

# Response Surface Methodology (RSM) for analysing culture conditions of *Acidocella facilis* strain USBA-GBX-505 and Partial Purification and Biochemical Characterization of Lipase 505 LIP

Luisa Fernanda Bernal<sup>1,2</sup>, Gina López<sup>1,2</sup>, Mónica Ruiz<sup>2</sup>, Ricardo Vera-Bravo<sup>3</sup>,  
Alejandro Reyes<sup>3</sup>, Sandra Baena<sup>1,2,\*</sup>

## Edited by

Juan Carlos Salcedo-Reyes  
(salcedo.juan@javeriana.edu.co)

1. Colombian Centre for Genomics and Bioinformatics of Extreme Environments - GeBiX.  
Carrera 5 No. 66A-34, 110231,  
Bogotá D.C. - Colombia.

2. Unidad de Saneamiento y Biotecnología Ambiental (USBA),  
Departamento de Biología, Facultad de Ciencias, Pontificia Universidad Javeriana, POB 56710,  
Bogotá D.C. - Colombia.

3. Química de Macromoléculas.  
Facultad de Ciencias, Pontificia Universidad Javeriana. POB 56710,  
Bogotá D.C. - Colombia

\* baena@javeriana.edu.co

Received: 04-02-2016

Accepted: 28-12-2016

Published on line: 16-02-2017

**Citation:** Bernal LF, López G, Ruiz R, Vera-Bravo R, Reyes A, Baena S. Response Surface Methodology (RSM) for analysing culture conditions of *Acidocella facilis* strain USBA-GBX-505 and Partial Purification and Biochemical Characterization of Lipase 505 LIP, *Universitas Scientiarum*, 22 (1): 45-70, 2017. doi: 10.11144/Javeriana.SC22-1.rsmr

## Abstract

Using Response Surface Methodology (RSM) we evaluated the culture conditions (nitrogen source, carbon source, pH and agitation rate) that increase the biomass of *Acidocella facilis* strain USBA-GBX-505 and therefore enhance the production of its lipolytic enzyme, 505 LIP. RSM results revealed that yeast extract and agitation were key culture factors that increased the growth-associated lipolytic activity by 4.5-fold (from 0.13 U.mg<sup>-1</sup> to 0.6 U.mg<sup>-1</sup>). The 505 LIP lipase was partially purified using size-exclusion chromatography and ion-exchange chromatography. Its molecular weight was >77 kDa. The enzyme shows its optimum catalytic activity at 55 °C and pH 7.5. EDTA, PMSF, 1-butanol and DMSO inhibited enzymatic activity, whereas Tween 20, acetone, glycerol and methanol increased it. Metallic ions are not required for the activity of 505 LIP, and even have an inhibitory effect on the enzyme. This study shows the potential use of *A. facilis* strain USBA-GBX-505 for the production of a newly identified lipolytic enzyme, 505 LIP, which is stable at moderate temperatures and in the presence of organic solvents. These are important characteristics for the synthesis of many useful products.

**Keywords:** lipases; acidophiles; *Acidocella facilis*; stability in solvents; Plackett-Burman design; central composite design.

## Introduction

Currently, 75 % of the enzymes used on an industrial scale are hydrolytic enzymes, with the carbohydrases, proteases and lipases dominating the enzyme market [1]. Lipases (EC 3.1.1.1) and esterases (EC 3.1.1.3) belong to the family of  $\alpha/\beta$  hydrolases, which are characterized by their capacity to hydrolyse and to form ester bonds in organic molecules, such as lipids and fats [1-3]. These enzymes are industrially important due to their capacity to act in environments with a low concentration of water, allowing them to perform esterification, transesterification, aminolytic and acidolytic reactions [4] with high efficiency and stability. The lipases are able to



**Funding:** This research was carried out with financial support from Pontificia Universidad Javeriana (Internal project: Identification of lipolytic enzymes produced by thermoacidophilic strains, partial purification and preliminary biochemical characterization ID 4269) and Colciencias – Sena. (Project 6570-392-199990).

**Electronic supplementary material:** Suppl.1

catalyse enzymatic reactions with efficiency, stability and chemo-, region- and enantio-selectivity, as well as not requiring cofactors and high activity levels in organic solvents make them versatile biocatalysers for industrial purposes [5], including food, dairy, agrochemical, pharmaceutical, cosmetic, and detergent production [5, 6] due to their unique properties, associated with their broad substrate range [7].

By 2017, the industrial demand for lipolytic enzymes will rise 6.2% [5], which means that it is necessary to continue to research and develop technology relating to them. Many of the lipolytic enzymes have poor stability during industrial processes, are denatured at high temperatures or extreme pH (alkaline or acid) and tend to lose activity over time at room temperature, resulting in inefficient catalysts. The industrial demand for lipolytic enzymes that are active at moderate and high temperatures and extreme pH has particularly encouraged the search for extremophilic microorganisms as source of thermostable enzymes [8-10]. Extremophilic microorganisms are the best source of extremozymes that catalyse reactions at extreme temperatures, extreme pH (acidic or alkaline pH), in the presence of solvents, high salinity, high radiation levels, low water activity or high concentrations of metals, among other environmental conditions [11]. However, some of these organisms exhibit slow and scarce growth, hindering the production of enzymes and therefore their subsequent purification and application. Identifying the culture variables that optimize the growth of the organisms that produce the enzymes is an important step in their production [12, 13]. To improve the microbial enzymatic production, it is necessary to consider several factors such as nutrients (sources of carbon and nitrogen), metal ions, pH, temperature, inoculum volume, among others [14].

The microbial communities of acidic environments are characterized by having very diverse and active metabolisms [15], mainly autotrophic and/or mixotrophic. These microorganisms produce active metabolites, including sesquiterpenoids, polyketide-terpenoid metabolites, berkeleydione, and berkeleytrione [16, 17]. They also produce stable extremozymes in conditions of extreme temperature and pH and are tolerant to solvents and detergents. Therefore, the search for extremozymes produced by acidophilic microorganisms represents a major challenge, and the exploration of pristine habitats offers possibilities to find enzymes with previously unreported features. The acidophilic organisms have been investigated not only for possible connections to ancient organisms but also for particular physical, structural, and molecular properties that allow them to live in acidic environments [18, 19]. During our previous studies on microbial diversity of acid hot springs in Los Nevados National Natural Park (Los Nevados NNP), we isolated several thermo and mesophilic acidophilic organisms with lipolytic activity tested on different substrates.

We evaluated the strain USBA-GBX-505 (CMPUJ U505 Acidocella, WDCM857), previously isolated from the acidic hot spring “El Coquito”(A4) (Los Nevados NNP) located at 3,973 m.a.s.l (4° 52' 41.99" N; 75° 16' 23.41" W) [20], as a source of lipolytic enzymes because our previous studies demonstrated that it is capable of hydrolyzing lipid substrates at ~55 °C. Additionally, we observed the highest lipolytic activity in the intracellular fraction compared to the extracellular fraction, but the strain exhibited a slow growth rate. In the present study we identified the growth factors of *A. facilis* strain USBA-GBX-505 that allow a higher production of biomass, which facilitated the partial purification and biochemical characterization of the lipolytic enzyme 505 LIP.

## Materials and Methods

### *Microorganism and culture conditions*

The acidophilic strain USBA-GBX-505 [20] was cultured in modified basal salt medium 991 (M991) [21] containing the following: (g.L<sup>-1</sup>) (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 1.3, KH<sub>2</sub>PO<sub>4</sub> 0.28, MgSO<sub>4</sub> · 7H<sub>2</sub>O 0.25, CaCl<sub>2</sub> · 2H<sub>2</sub>O 0.07, and FeSO<sub>4</sub> 0.02, supplemented with 1 g.L<sup>-1</sup> yeast extract, 5 g.L<sup>-1</sup> peptone and 10 ml.L<sup>-1</sup> trace element solution SL-10 [22]. The pH was adjusted to 3.5 by using 5 N H<sub>2</sub>SO<sub>4</sub>. The microorganism was incubated at 30 °C with agitation at 200 r/min. The inoculum used for all assays was standardized at an absorbance of 0.3 - 0.4 at 580 nm, corresponding to a concentration of ~4 x 10<sup>8</sup> CFU.mL<sup>-1</sup>, obtained during the mid-exponential phase of growth (48 hours).

### *Extraction of intracellular proteins*

The biomass was concentrated by centrifugation at 8,000 *g* for 20 min at 10 °C. The cells were washed twice with Tris-HCl buffer (50 mM, pH 7.5). Then, the biomass was resuspended in the same buffer with 200 µg.mL<sup>-1</sup> of lysozyme (Sigma-Aldrich, U.S.A.), and the mixture was incubated at 37 °C for 10 min and placed on ice for 30 min. Cell lysis was performed through sonication using a QSonica Q125 sonicator (Qsonica, Newtown, CT, U.S.A.), in 15 cycles of 30 s sonication pulses at 30% amplitude and 30 s breaks at 0 °C. Finally, the lysate was centrifuged at 8,000 *g* for 25 min at 4 °C, and the supernatant was considered to be the intracellular protein extract [20].

### *Evaluation of the intracellular lipolytic activity using p-nitrophenyl ester (pNP-ester)*

The lipolytic activity was measured spectrophotometrically using the pNP- decanoate (pNP-C10) as a substrate. The enzymatic reaction was performed under the following conditions: 425 µl of Tris-HCl buffer (50 mM, pH 7.5), 25 µl of the substrate pNP-ester at a final concentration of 0.5 mM and 25 µl of the sample [23]. The absorbance at 405 nm was measured using a HACH DR 5000™ UV-Vis Spectrophotometer (HACH Loveland, CO, U.S.A.). The blank was Tris-HCl buffer (50 mm, pH 7.5) containing the substrate pNP-C10 (0.5 mM). All of the measurements were performed by triplicate. The molar extinction coefficient used was 1.336 x 10<sup>7</sup> cm<sup>2</sup>. mol<sup>-1</sup>.

The specific lipolytic unit was defined as the amount of enzyme that catalysed the formation of 1 µmol of p-nitrophenol (pNP) per min per 1 mg of protein (U.mg<sup>-1</sup>). Quantification of the total protein was performed using the Bradford technique, with spectrophotometry at 595 nm [24] using bovine serum albumin as the standard.

### *Identification of culture variables for A. facilis strain USBA-GBX-505*

We used a Plackett-Burman (PB) fractional factorial design in order to identify the culture variables that favoured higher levels of biomass production and therefore enzymatic activity. The response surface methodology (RSM) was employed for obtaining the optimum concentration of individual variables that potentially influence the response (lipolytic activity) [25]. The PB included eight culture factors at two levels, high (+1) and low (-1): peptone and yeast extract (5 and 10 g.L<sup>-1</sup>) as the organic nitrogen source; ammonium chloride (NH<sub>4</sub>Cl, 0.1 and 1.0 g.L<sup>-1</sup>) and ammonium sulfate ((NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0 and 1.0 g.L<sup>-1</sup>) as the inorganic nitrogen source; Tween 80 and

ethyl oleate (0.1 and 1% (v/v)) as the carbon source and inducer of lipolytic activity; pH (3.0 and 5.0); and agitation (0 and 200 r/min). Each experimental round was conducted by triplicate. The dependent variable was the specific intracellular lipolytic activity ( $\text{U}\cdot\text{mg}^{-1}$ ) obtained from the cultured biomass after 96 h of incubation.

We used Design-Expert (version 8) software (Stat-Ease, Inc. U.S.A.) to determine the level of experimental error, corroborate the fit of the first-order model and perform regression analysis of the experimental results obtained.

### *Central composite design (CCD)*

The experimental region of the CCD comprised five levels (-1.68, -1, 0, 1, and 1.68) of each of the factors under evaluation [26]. The levels of yeast extract, Tween 80<sup>®</sup> and agitation were evaluated in twenty treatments using Design-Expert (version 8) software, generating a CCD matrix that included axial and central points. All of the treatments were evaluated by triplicate. The dependent variable was the specific intracellular lipolytic activity ( $\text{U}\cdot\text{mg}^{-1}$ ).

### *Statistical analysis of experimental designs*

The quality of fit obtained using the first-order-model equation was expressed as the determination coefficient  $R^2$ ; its significance was determined using an  $F$  (Fisher) test, and the statistical significance of the regression coefficients was corroborated using the Student's  $t$  test. Analyses of variance were performed to evaluate the results obtained using the PB design, CDD and quadratic model of cultures of *A. facilis* USBA-GBX-505 to be used for the production of lipolytic enzyme. Data obtained from the RSM on lipase production were subjected to analysis of variance (ANOVA).

### *Partial purification of lipase 505 LIP*

We used the following culture conditions for the production of USBA-GBX-505 microbial biomass: modified basal salt 991 medium, supplemented with 6.6  $\text{g}\cdot\text{L}^{-1}$  of yeast extract and 1% (v/v) Tween 80<sup>®</sup>, and incubation at 30 °C for 4 d, with agitation at 200 r/min. The intracellular protein obtained as described above, was concentrated and pre-purified using an Amicon ultrafiltration unit of 50 kDa (Merck Millipore Corporation, Darmstadt, Germany) with centrifugation at 3,000  $g$  for 15 min. The resulting extract was loaded on a BioSep-SEC-3000 size-exclusion column (Shodex<sup>®</sup>, Germany) of 30 cm x 0.78 cm that had been equilibrated with 50 mM Tris-HCl buffer (pH 7.5) and was subsequently eluted at a flow rate of 1  $\text{mL}\cdot\text{min}^{-1}$  at 35 °C for 20 min, with detection at 215 nm. We collected and then concentrated by ultrafiltration fractions of 2.5 mL for subsequent protein quantification, evaluation of enzymatic activity and enzyme purification. The fraction exhibiting enzymatic activity that was collected using size-exclusion chromatography (SEC) was loaded on an IEC-DEAE-2025 ion-exchange column (Shodex<sup>®</sup>) of 15 cm x 2 cm. Then, 50 mM Tris-HCl buffer, pH 7.5 was applied at a flow rate of 1  $\text{mL}\cdot\text{min}^{-1}$  at 35 °C for 20 min, after which a gradient of 0 to 1 M NaCl in 50 mM Tris-HCl, pH 7.5 was applied between 20 and 110 min, with detection at 280 nm. We collected, desalted and concentrated by ultrafiltration the subsequent fractions for the quantification of the protein and lipolytic enzymatic activity. The fractions obtained using SEC and ion-exchange chromatography (IEC) were evaluated using a diode array detector.

The resulting protein fractions were analysed by electrophoresis in a 10% (w/v) polyacrylamide gel containing sodium dodecyl sulfate (SDS-PAGE) using a Mini-Protean II<sup>®</sup> system (BioRad, Bio-Rad Laboratories, Inc, U.S.A.) according to the Laemmli protocol [27]. Gel staining was performed using an ammoniacal silver-staining protocol [28]. The protein molecular weight marker used was V849A (Promega, USA).

## Biochemical characterization of purified enzyme 505 LIP

For all the characterization assays, the quantification of the lipolytic activity of enzyme 505 LIP was performed using the *p*-nitrophenyl-ester technique, with 0.5 mM *p*NP decanoate (C10) as the substrate, except for substrate specificity evaluations. We calculated the lipolytic activity as the percentage of the residual activity, with 100% activity corresponding to the control (the enzyme without any interfering compound). The evaluations were performed using *p*NP-C10 at 55 °C in Tris-HCl buffer (50 mM, pH 7.5).

### *Substrate specificity evaluations*

#### Spectrophotometric method

The evaluation was done using *p*-NP ester substrates (Sigma-Aldrich) (described above) with different chain lengths at 50 °C and Tris-HCl buffer (50 mM, pH 7.5). The *p*-NP esters evaluated were acetate (C2), butyrate (C4), octanoate (C8), decanoate (C10), dodecanoate (C12), hexadecanoate (C16) and octadecanoate (C18). All *p*-nitrophenyl esters were dissolved in 2-propanol to a final concentration of 0.5 mM. The *p*-nitrophenyl esters > C10 were also mixed with Triton X-100 (0.002 % (w/v) final concentration) (Sigma-Aldrich).

#### Titrimetric method

We used the titration method to determine the free fatty acids concentrations, according to Seghal et al. (2014) [7]. Briefly, the reaction mixture contained 5% (w/v) of the lipidic substrate (tributyrate, tricaproline and trioleate) in Tris-HCl buffer (50 mM, pH 7.5) and 1% of Triton X-100. The reactions were incubated at 50 °C and 200 r/min for 30 min. Then, we added acetone-ethanol mixture (1:1 v/v). The final analysis was carried out using a potentiometric titration technique with 50 mM KOH (end point pH 8.0).

### *Evaluation of the optimal temperature and pH for enzymatic activity*

We evaluated the enzymatic activity at different temperatures in order to determine the optimal one, by exposing the enzyme for 30 min to temperatures ranging between 30 °C and 90 °C in Tris-HCl buffer (50 mM, pH 7.5). Then, the lipolytic activity was quantified as described above. To evaluate the enzymatic activity under different pH conditions, different buffer solutions were used at 50 mM, including MES (pH 5.2 – 6.7), Tris HCl (pH 7.0 – 9.0), HEPES (pH 6.8 – 8.2), BICINE (pH 7.6 – 9.0) and sodium carbonate (pH 9.0 – 11.0). The enzyme was incubated for 30 min at room temperature (23 °C ± 2) and then the lipolytic activity was quantified at 55 °C using *p*NP-C10.

### *Effect of various additives on the activity of 505 LIP*

We evaluated the effect of different chemical substances on the enzymatic activity by incubating the protein in the presence of each substance for 1 h at room temperature ( $23\text{ }^{\circ}\text{C} \pm 2$ ). We evaluated the effect of metallic ions as  $\text{Ca}^{2+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Cu}^{2+}$ , and  $\text{Mn}^{2+}$  at two concentrations, 1 mM and 10 mM, except for the  $\text{Fe}^{3+}$  ion, which was evaluated at 1 mM and 5 mM due to the interference caused by the colour formation at 10 mM. The effect of sodium dodecyl sulfate (SDS), Tween 20, Tween 80 and Triton X-100 was evaluated at concentrations of 1 and 5% (v/v). The effect of the inhibitors phenylmethylsulfonyl fluoride (PMSF), ethylenediaminetetraacetic acid (EDTA), 2-mercaptoethanol (2-ME), and dithiothreitol (DTT) was evaluated at 5 mM as final concentration.

### *Effect of organic solvents on the stability of 505 LIP*

The enzymatic activity was also evaluated in the presence of several organic solvents, including 1-butanol, acetonitrile, acetone, 1-propanol, 2-propanol, methanol, ethanol, glycerol and dimethyl sulfoxide (DMSO) at 15 and 30% (v/v). The stability of the enzyme was determined in the presence of 15% (v/v) methanol and 30% (v/v) glycerol at room temperature ( $23\text{ }^{\circ}\text{C} \pm 2$ ) for 192 h. The reaction conditions were  $55\text{ }^{\circ}\text{C}$  and pH 7.5, and the substrate used was *p*-NPC10 at 0.5 mM.

All of the chemical products used were of the purest grades available.

## **Results and Discussion**

While searching for extremozymes produced by thermophilic and acidophilic microorganisms in the acidic hot springs at Los Nevados NNP, we isolated several microorganisms with diverse characteristics and lipolytic activities such as the thermoacidophilic alpha-Proteobacterium *Acidicaldus* USBA-GBX-499 [15] and the *A. facilis* strain USBA-GBX-505 [20].

In order to identify the factors that significantly affect the growth of *A. facilis* USBA-GBX-505 for the production of biomass and lipolytic enzymes, a PB fractional factorial design was developed. Treatment 2 exhibited the highest lipolytic activity ( $0.23\text{ U}\cdot\text{mg}^{-1}$ ). This treatment consisted of the organic nitrogen sources (peptone and yeast extract) and ethyl oleate at a high level ( $10\text{ g}\cdot\text{L}^{-1}$  and 1% v/v), the sources of inorganic nitrogen ( $\text{NH}_4\text{Cl}$  and  $(\text{NH}_4)_2\text{SO}_4$ ) at a low level ( $0.1\text{ g}\cdot\text{L}^{-1}$  and  $0\text{ g}\cdot\text{L}^{-1}$ , respectively), and the cultures incubated at high level agitation rate (200 r/min). In contrast, the lowest level of activity was detected in treatment 8 ( $0.002\text{ U}\cdot\text{mg}^{-1}$ ) and no lipolytic activity was detected ( $0\text{ U}\cdot\text{mg}^{-1}$ ) in treatments 10 and 12. All three of these treatments had low levels of agitation (0 r/min) and yeast extract ( $5\text{ g}\cdot\text{L}^{-1}$ ). These results showed that little or no growth of the microorganism occurred in the absence of agitation or with a low level of yeast extract.

The analysis of variance (ANOVA) of the PB design (**Table 1**) showed that the variables yeast extract, Tween 80<sup>®</sup> and agitation were the culture factors that significantly ( $p < 0.05$ ) affected the production of biomass and hence of lipolytic enzymes. The nitrogen source for any microorganism serves as a secondary energy source for growth and enzyme secretion; the nitrogen is especially important in

**Table 1.** Matrix and analysis of variance (ANOVA) of the Plackett-Burman experimental design used to evaluate various culture factors of *Acidocella facilis* USBA-GBX-505 for the production of lipolytic enzymes. T80: Tween 80®, EO: Ethyl oleate, Pep: Peptone, YE: Yeast extract, D<sub>1</sub>: Dummy1, D<sub>2</sub>: Dummy2, D<sub>3</sub>: Dummy3, TTo: Treatment. \*Specific intracellular lipolytic activity after 48 h of culture at 30 °C.

TTo	A	B	C	D	E	F	G	H	I	J	K	* specific lipolytic activity (U.mg <sup>-1</sup> )	
	T80 %(v/v)	EO %(v/v)	Pep (g/l)	YE (g/l)	(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> (g/l)	NH <sub>4</sub> Cl (g/l)	pH	Agitation (rpm)	D <sub>1</sub>	D <sub>2</sub>	D <sub>3</sub>	Experimental	Prediction
1	+1	-1	+1	+1	+1	-1	-1	-1	1	1	-1	0.027	0.033
2	-1	+1	+1	+1	-1	-1	-1	+1	-1	1	1	0.229	0.161
3	-1	-1	-1	+1	-1	+1	+1	-1	1	1	-1	0.012	0.018
4	+1	-1	-1	-1	+1	-1	+1	+1	-1	1	1	0.177	0.151
5	-1	-1	+1	-1	+1	+1	-1	+1	1	-1	-1	0.027	0.037
6	+1	-1	+1	+1	-1	+1	+1	+1	-1	-1	1	0.214	0.171
7	+1	+1	-1	+1	+1	+1	-1	-1	-1	1	-1	0.182	0.184
8	-1	+1	+1	-1	+1	+1	+1	-1	-1	-1	1	0.002	0.004
9	+1	+1	-1	-1	-1	+1	-1	+1	1	1	1	0.022	0.034
10	+1	+1	+1	-1	-1	-1	+1	-1	1	-1	1	0.000	0.016
11	-1	+1	-1	+1	+1	-1	+1	-1	1	-1	-1	0.095	0.099
12	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	0.000	0.003
<b>SS</b>	5.704 x 10 <sup>-4</sup>	4.619 x 10 <sup>-4</sup>	1.524 x 10 <sup>-4</sup>	7.168 x 10 <sup>-3</sup>	1.251 x 10 <sup>-4</sup>	4.445 x 10 <sup>-4</sup>	9.702 x 10 <sup>-5</sup>	7.418 x 10 <sup>-3</sup>	2.340 x 10 <sup>-3</sup>				
<b>df</b>	1	1	1	1	1	1	1	1	1	1	1		
<b>RMS</b>	5.704 x 10 <sup>-4</sup>	4.619 x 10 <sup>-4</sup>	1.524 x 10 <sup>-4</sup>	7.168 x 10 <sup>-3</sup>	1.251 x 10 <sup>-4</sup>	4.445 x 10 <sup>-4</sup>	9.702 x 10 <sup>-5</sup>	7.418 x 10 <sup>-3</sup>	2.340 x 10 <sup>-3</sup>				
<b>F Value</b>	20.87	16.90	5.58	262.30	4.98	16.27	3.55	271.44	85.64				
<b>P Value</b>	0.0447*	0.0544	0.1421	0.0038*	0.0613	0.0563	0.2002	0.0037*	0.0115				

regulating the synthesis of hydrolases [29]. Sharma et al., (2012) [30] showed that yeast extract was the main nitrogen source for lipase production by *B. licheniformis* MTCC-10498. In contrast, ammonium phosphate was the main source of nitrogen for lipase production by *G. thermoleovorans* DA2 [31]. The Tween 80® is widely used as a carbon source by microorganisms producing esterases such as *Acidiphilum* sp. AIU (formally, *Acidocella facilis*) [32] and *Acidocella aminolytica* [33]. From our results, ethyl oleate, peptone, inorganic nitrogen sources ((NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, NH<sub>4</sub>Cl) and pH did not have a significant effect.

The first-order model generated through the data analysis indicated a sample variation of 98.9%. Therefore, we can deduce that there was a relationship between the culture factors of *A. facilis* USBA-GBX-505 and its production of lipolytic enzyme. Only 1.1% of the total variation could not be explained by the model. Even the R<sup>2</sup><sub>Pred</sub> of 0.929 demonstrated that there was a strong correlation between the experimental values and the values predicted using the following linear model:

#### Equation 1. First order model:

$$\text{U.mg}^{-1} = 0.038 + 6.9 \times 10^{-3}\mathbf{A} + 6.2 \times 10^{-3}\mathbf{B} + 3.5 \times 10^{-3}\mathbf{C} + 0.024\mathbf{D} - 0.028\mathbf{E} - 6.1 \times 10^{-3}\mathbf{F} - 2.8 \times 10^{-3}\mathbf{G} + 0.025\mathbf{H} + 0.014\mathbf{K} \quad (1)$$

where **A**: Tween80®, **B**: Ethyl oleate, **C**: Peptone, **D**: Yeast extract, **E**: (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, **F**: NH<sub>4</sub>Cl, **G**: pH, **H**: Agitation, and **K**: Dummy2.

The positive regression coefficient of 0.024 ( $p = 0.0038$ ), 0.025 ( $p = 0.0037$ ) and  $6.9 \times 10^{-3}$  ( $p=0.0447$ ) for yeast extract, agitation and Tween 80®, respectively, confirmed that these factors had a significantly positive effect on the production of lipolytic enzymes by *A. facilis* USBA-GBX-505. Therefore, we used these factors in the CCD experiment.

Considering that all three significant factors had a positive effect in the PB-design-based experiment, we decided that the highest levels of yeast extract and Tween 80®, previously evaluated using the PB design, would be the low levels in the CCD-based experiment. Hence, in the CCD-based experiment, the low level of yeast extract was 10 g.L<sup>-1</sup> and that of Tween 80® was 1 % (v/v). In the PB-design-based experiment, two agitation levels were evaluated, low (0 r/min) and high (200 r/min), and a tendency towards a higher lipolytic activity was observed in the treatments with a high agitation level; therefore we decided to evaluate agitation rates that were lower and higher than 200 r/min. Three factors were evaluated at 5 levels for a total of 20 treatments. **Table 2** shows the composition of each of the experimental treatments and their respective specific lipolytic activities.

Treatment 9 had the maximal intracellular lipolytic activity of 0.6 U.mg<sup>-1</sup> experimental and 0.5 U.mg<sup>-1</sup> predicted, as well as the highest biomass level of  $8 \times 10^7$  CFU.mL<sup>-1</sup>. In contrast, treatments 2, 6, 8, 10 and 12 did not show any lipolytic activity (Table 2). **Table S1** shows the ANOVA results used to compare the mean values at  $p = 0.001$  ( $p < 0.05$ ), which demonstrated that there were significant differences among the treatments. The results confirmed that treatment 9 independently stimulated the production of lipases by *A. facilis* USBA-GBX-505 and that the levels of Tween 80®, yeast extract and agitation used in treatment 9 enhanced the production of lipolytic



**Table 2.** Matrix of the experimental central composite design (CCD) used to evaluate 3 culture factors of *Acidocella facilis* USBA-GBX-505 for the production of lipolytic enzymes. YE: Yeast extract, TTo: Treatment. \*Specific intracellular lipolytic activity after 48 h of culture at 30 °C.

TTo	A	B	C	*Specific lipolytic activity (U.mg <sup>-1</sup> )	
	YE (g.L <sup>-1</sup> )	Agitation (r/min)	T80 %(v/v)	Experimental	Prediction
1	-1(10)	-1(150)	-1(1)	0.237	0.231
2	+1(20)	-1(150)	-1(1)	0.000	0.040
3	-1(10)	+1(250)	-1(1)	0.042	0.049
4	+1(20)	+1(250)	-1(1)	0.047	0.048
5	-1(10)	-1(150)	+1(2)	0.074	0.094
6	+1(20)	-1(150)	+1(2)	0.000	0.045
7	-1(10)	+1(250)	+1(2)	0.163	0.164
8	+1(20)	+1(250)	+1(2)	0.000	0.025
9	-1.68(6.6)	0(200)	0(1.5)	0.590	0.495
10	1.68(23.4)	0(200)	0(1.5)	0.000	0.059
11	0(15)	-1.68(116)	0(1.5)	0.052	0.061
12	0(15)	1.68(284)	0(1.5)	0,000	0.018
13	0(15)	0(200)	-1.68(0.7)	0.207	0.206
14	0(15)	0(200)	1.68(2.3)	0.123	0.122
15	0(15)	0(200)	0(1.5)	0.248	0.240
16	0(15)	0(200)	0(1.5)	0.245	0.240
17	0(15)	0(200)	0(1.5)	0.244	0.240
18	0(15)	0(200)	0(1.5)	0.245	0.240
19	0(15)	0(200)	0(1.5)	0.245	0.240
20	0(15)	0(200)	0(1.5)	0.244	0.240

enzymes by this chemoorganotrophic organism whose growth is favoured by protein substrates [34]. These substrates most likely served not only as nitrogen sources but also as sources of B-complex vitamins, as reported before [35]. Moreover, previous reports indicate that the sources of organic nitrogen affect the growth of the organism and are directly related to the production of lipases [36].

The experimental data obtained were analysed using the surface response methodology, and the fit of the second-order model was evaluated and the coefficients of the second-order polynomial were generated through multiple regressions. The quality of the fit of the model ( $p = 0.0433$ ) was evaluated using the determination coefficients and an ANOVA. The quadratic model of the response surface was adjusted using the following equation:

**Equation 2. Second-order model:**

$$U.mg^{-1} = 0.24 - 0.10A - 0.013B - 0.025C + 0.025 AB + 5.5 \times 10^{-3}AC + 0.023BC - 2.5 \times 10^{-3} A^2 - 0.099B^2 - 0.027C^2 \quad (2)$$

where: **A:** Yeast extract, **B:** Agitation, and **C:** Tween 80®.


Based on the results of the ANOVA of the CCD-based experiment (**Table 3**), the Fisher F value at 3.17 and the p value (0.0433,  $p < 0.05$ ), we consider that the model (**Equation 2**) was significant. The highest F values were observed for yeast extract (A) and agitation2 (B2); therefore, these two values had a significant effect ( $p < 0.05$ ) on the growth of *A. facilis* USBA-GBX-505 for the production of lipolytic enzymes. Likewise, as shown in Table 3, the quadratic effect of the agitation rate ( $p = 0.0049$ ) was significant compared with its linear effect ( $p = 0.6560$ ). The other significant value for the production of lipases was the linear effect of the yeast extract level ( $p = 0.0046$ ). In summary, yeast extract and agitation were the factors that had a significant effect ( $p < 0.05$ ) on the growth of *A. facilis* USBA-GBX-505 for the production of lipolytic

**Table 3.** Analysis of variance (ANOVA) of the data obtained using the CCD-based quadratic model of *Acidocella facilis* USBA-GBX-505. SS: Sum of squares, df: Degrees of freedom, RMS: Root mean square  $R^2$ : 0.7954,  $R^2_{Adjust}$ : 0.5970, Adeq Precision: 6.932, CV%: 9.69, Std Dev: 0.10 \*Significance at  $p < 0.05$ .

Variable	SS	df	RMS	F Value	P Value
<b>Model</b>	0.031	9	0.035	3.17	0.0433*
<b>A-YE</b>	0.15	1	0.15	13.23	0.0046*
<b>B-Agitation</b>	$2.321 \times 10^{-3}$	1	$2.321 \times 10^{-3}$	0.21	0.6560
<b>C-T80</b>	$8.861 \times 10^{-3}$	1	$8.861 \times 10^{-3}$	0.81	0.3908
<b>AB</b>	$5.109 \times 10^{-3}$	1	$5.109 \times 10^{-3}$	0.46	0.5113
<b>AC</b>	$2.459 \times 10^{-4}$	1	$2.459 \times 10^{-4}$	0.022	0.8842
<b>BC</b>	$4.368 \times 10^{-3}$	1	$4.368 \times 10^{-3}$	0.40	0.5429
<b>A<sup>2</sup></b>	$8.651 \times 10^{-5}$	1	$8.651 \times 10^{-5}$	$7.855 \times 10^{-3}$	0.9311
<b>B<sup>2</sup></b>	0.15	1	0.15	12.91	0.0049*
<b>C<sup>2</sup></b>	0.011	1	0.011	0.96	0.3498
<b>Residual</b>	0.11	10	0.11		

enzymes. **Figure 1** shows the interaction between these two factors in which the best response was observed using an agitation rate of 200 r/min and 10 g.L<sup>-1</sup> of yeast extract. An inversely proportional linear effect of the yeast extract level and the specific lipolytic activity was also observed. This result indicated the possibility that at a yeast extract concentration of less than 10 g.L<sup>-1</sup>, lipase production might increase. This result is similar to that reported for *Burkholderia* sp. strain C20, for which a concentration of yeast extract greater than 2 g.L<sup>-1</sup> caused a decrease in lipase production, likely due to substrate inhibition of the nitrogen source [37]. The other significant factor for the production of lipolytic enzymes by *A. facilis* USBA-GBX-505 was the agitation rate. The agitation rate is one of the most important culture parameters because increasing this rate increases the oxygen transfer rate as well as the mixing efficiency of the culture, finally leading to increased cellular growth and lipase production [38]. Such is the case of *Acinetobacter radioresistens*, for which the optimal yield of lipases was observed at an agitation rate of 600 r/min [39].

In summary, we observed the highest specific activity in treatment 9, which increased the enzymatic activity level by 4.5-fold compared with that obtained using the initial culture conditions (agitation at 0 r/min, 5 g.L<sup>-1</sup> peptone, 1 g.L<sup>-1</sup> yeast extract and 1% (v/v) Tween 80<sup>®</sup>), under which the intracellular specific enzymatic activity was only 0.13 U.mg<sup>-1</sup> (data not shown). Tween 80<sup>®</sup> is one of the factors improving the production of the lipase enzyme by *A. facilis* strain USBA-GBX-505 in the first screening experiments conducted with the PB-based design; it has also been reported to be important for



**Fig. 1.** Response surface curve showing the effect of yeast extract (YE) and agitation on the culture of *Acidocella facilis* USBA-GBX-505 for the production of lipolytic enzymes.


other organisms, such as *Burkholderia glumae* PG1 [40] and *Staphylococcus epidermidis* AT2 [41]. Tween 80<sup>®</sup>, a surfactant agent, is also used as a substrate for the production of lipases and esterases, in many cases efficiently increasing their synthesis rate [42]. For *A. facilis* USBA-GBX-505, this compound could serve as the only carbon source and as an inducer of lipolytic activity.

The analysis of the factors associated with the production of microbial lipolytic enzymes has been the focus of various studies because it is known that each microorganism has specific requirements and particular conditions for the maximization of the production of enzymes. Among the factors associated with the production of lipases are lipid inducers, which are mainly used as carbon sources, organic and inorganic nitrogen sources and oligo elements, among others [36]. To date, the only knowledge regarding optimization of the production of lipolytic enzymes by an acidophilic microorganism concerned a strain of *Bacillus pumilus*, which grows at pH 1.0, for which the production of an acid lipase was enhanced by optimizing the pH, temperature and lipid-substrate concentration (palm oil: 5% (v/v)) [43]. Likewise, there are reports concerning fungi such as *Aspergillus wentii* that grow at a pH lower than 5.0, for which the yield of lipase was increased by adding peptone and yeast extract to the culture medium [44].

Our results demonstrated the culture conditions for *A. facilis* strain USBA-GBX-505 that increased its lipase production, which were as follows: modified 991 medium supplemented with yeast extract at 6.6 g·L<sup>-1</sup> and Tween 80<sup>®</sup> at 1.5% (v/v), with an agitation rate of 200 r/min. Next, we proceeded to culture *A. facilis* USBA-GBX-505 under these particular conditions to obtain a protein extract with lipolytic activity and then to biochemically characterize the lipolytic enzyme it produces.

#### *Partial purification of the lipolytic enzyme of A. facilis USBA-GBX-505*

We established a purification method for lipolytic enzymes based on semi-preparative high-performance liquid chromatography (HPLC). The intracellular protein extract, obtained as previously described, was concentrated using 50 kDa Amicon ultrafiltration units, yielding two fractions: the first fraction containing proteins of < 50 kDa and the second fraction containing proteins of > 50 kDa. Assessing the lipolytic activity of the two fractions demonstrated that the second fraction had lipolytic activity. Therefore, we used this fraction for subsequent purification using SEC, from which three fractions were collected; the first fraction, collected between 5 and 7.5 min, showed lipolytic activity. This fraction was concentrated using a filtration system designed for 50 kDa proteins, after which IEC was used to increase the purity level of the active fraction. This fraction was eluted from the ion-exchange column at 180 min in 0.8 M NaCl/50 mM Tris-HCl (pH 7.5). The changes in the purity level of the active fraction throughout the partial purification process and the molecular mass of the enzyme were determined using SDS-PAGE (**Figure 2**). In the third lane of the SDS-PAGE gel shown in **Figure 2**, which corresponds to the active fraction obtained using IEC, multiple bands were observed, indicating that the active protein was partially purified. This enzyme, denominated 505 LIP, had a molecular weight greater than 77 kDa. The extracellular esterase that was isolated and purified from *A. facilis* strain AIU409 using genomic methods had a molecular weight of ~64 kDa [45], indicating that that esterase and the 505 LIP are two different kinds of esterases, which are expressed in two different strains of the same species.




**Fig. 2.** SDS - PAGE electrophoresis gel (10% (w/v)) of *Acidocella facilis* strain USBA-GBX-505 lipase purification process. Lane 1: Crude extract, Lane 2: Active fraction collected using size-exclusion chromatography (SEC), Lane 3: Active fraction collected using ion-exchange chromatography (IEC), Lane 4: 6.5-200 kDa V849A (Promega, USA).

The enzyme 505 LIP was purified with a recovery yield of 16%. The specific activity of this enzyme was  $6.3 \text{ U}\cdot\text{mg}^{-1}$ , and the recovery factor of its activity was 2.3 times greater than prior to purification (**Table 4**). Several authors have stated that protein purification yields of between 2 and 20 % could be due to the formation of protein aggregates during purification because a large amount of enzyme can be lost as insoluble material, particularly when solutions with a pH of  $< 12$  are used [46, 47].

**Table 4.** Results of the strategy used to purify the lipolytic enzyme 505 LIP of *Acidocella facilis* USBA-GBX-505. <sup>1</sup>Active fraction collected using a size-exclusion column (SEC), <sup>2</sup>Active fraction collected using an ion-exchange column (IEC).

Step of purification	Total proteins (mg)	Total Activity (U)	Specific activity ( $\text{U}\cdot\text{mg}^{-1}$ )	Recovery factor	Recovery Yield %
Intracellular concentrated proteins	44.5	125.67	2.8	1.0	100
SEC Fraction <sup>1</sup>	15.1	78.33	5.0	1.8	62
IEC Fraction <sup>2</sup> (Partial purification extract 505LIP)	3.25	20.54	6.3	2.3	16




**Fig. 3.** Evaluation of lipolytic activity of the enzyme 505 LIP on *p*-nitrophenyl (*p*NP) esters with fatty acid chains of 4 to 18 C-atoms. Activities are shown as percentages of the maximum activity (10.96 U/mg). Values are means  $\pm$  SD from three independent experiments.

### Biochemical characterization of the lipolytic enzyme 505 LIP

The highest enzymatic activity of the 505 LIP was exhibited towards medium-chained *p*NP esters (**Figure 3**), such as decanoate (C10) and dodecanoate (C12), a finding that allowed to establish that the 505 LIP is a lipase, given that true lipases are enzymes that hydrolyse long-chain acylglycerols (with > 10 carbon atoms) and natural lipids [3]. However, its activity towards C16- and C18-acylglycerols was greatly decreased and a low level of enzymatic activity towards short- (C4) and medium-chained (C8) substrates was observed. Moreover, we evaluated the lipolytic activity on tributyrine, tricapriline and trioleate, and the results showed partial hydrolysis of 5, 40 and 10 % respectively after 24 h of reaction.

The optimal temperature for the 505 LIP activity was measured in the range of 30 °C to 90 °C (**Figure 4**) using *p*NP-C10 as the substrate. The highest levels of activity were observed between 45 °C and 60 °C, with 55 °C being the temperature at which the maximal activity toward *p*NP-C10 was detected. The lipolytic activity decreased by more than 50% at temperatures of less than 40 °C and greater than 65 °C. Therefore, the enzyme 505 LIP is classified as moderately thermophilic, with an optimal lipolytic activity temperature of 55 °C. The optimal temperature of the enzyme 505 LIP also differed from that of the esterase produced by *A. facilis* AIU409, which had an optimal temperature of 70 °C [45]. However, the optimal temperature of the 505 LIP is similar to that of the lipase derived from the acidophilic organism *B. pumilus* (50 °C). Other lipolytic enzymes derived from acidophilic organisms include the lipolytic



**Fig. 4.** Evaluation of lipolytic activity of the enzyme 505 LIP on *p*-nitrophenyl (*p*NP) esters with fatty acid chains of 4 to 18 C-atoms. Activities are shown as percentages of the maximum activity (10.96 U/mg). Values are means  $\pm$  SD from three independent experiments.

enzyme produced by the thermoacidophilic microorganism *Alicyclobacillus acidocaldarius*, with an optimal temperature of 60 °C [48], and the thermophilic esterase produced by *Acidicaldus* sp. USBA-GBX-499, with an optimal temperature of 55 °C [23].

The lipolytic activity of the 505 LIP was detected at pH values of between 6.0 and 9.0, with maximal activity observed at pH 7.5 (9.76 U $\cdot$ mg<sup>-1</sup> lipolytic activity of 505 LIP), and the enzymatic activity decreased as the pH became more alkaline. In contrast, the optimal pH for the growth of *A. facilis* strain USBA-GBX-505 was pH 3.5. The optimal pH value for the growth of this strain is probably different to the intracellular pH values of several acidophilic organisms that have various strategies for adaptation under acidic pH conditions, which are related to maintaining their intracellular pH at an approximately neutral value [49]. This result contrasts with that reported for the extracellular esterase of *A. facilis* AIU409, whose optimal activity was observed at pH 4.0 [45]. This optimal pH also differs from that of the lipase produced by the acidophilic microorganism *B. pumilus*, which had an optimal enzymatic activity at pH 1.0. The enzyme activity also could be affected by the buffer composition, because the enzymes have preferences for ionic strength (salt concentration) and major cation (sodium or potassium).

#### *Effect of metallic ions on the activity of the enzyme 505 LIP*

The activity of the enzyme 505 LIP decreased in the presence of various ions at 1 and 10 mM (Table 5), by between 40 and 100%, becoming totally inhibited in the presence of Fe<sup>2+</sup> ions. This finding is contrary to those previously reported on lipolytic enzymes,

**Table 5.** Effect of different chemical compounds on the activity of the enzyme 505 LIP determined using *p*NP-C10 as the substrate at pH 7.5 and 55 °C. A total of 100% activity was set as that observed in the absence of the compound under evaluation. \*Fe<sup>3+</sup> was the only ion evaluated at 1 and 5 mM due to the interference caused by the color of the 10 mM ionic solution. 100% Activity = 6.3 U.mg<sup>-1</sup>. The assays were performed in triplicate, and the mean values and standard deviations are shown.

Metal ion	Residual Activity (%)		
	1 mM	5 mM	10 mM
Cd <sup>2+</sup>	69 ± 1.1	-	59 ± 1.42
Cu <sup>2+</sup>	58 ± 0.5	-	13 ± 0.61
Ni <sup>2+</sup>	62 ± 1.3	-	61 ± 0.30
Fe <sup>2+</sup>	0 ± 0	-	0 ± 0
*Fe <sup>3+</sup>	56 ± 0.26	71 ± 1.06	-
Organic Solvent	15%	30%	
1-Butanol	0 ± 0	0 ± 0	
1-Propanol	47 ± 0.79	7 ± 0.40	
2-Propanol	53 ± 0.13	6 ± 0.34	
Acetone	114 ± 1.55	34 ± 1.52	
Ethanol	66 ± 0.35	32 ± 0.08	
Methanol	123 ± 0.86	88 ± 0.04	
DMSO	0 ± 0	0 ± 0	
Acetonitrile	62 ± 0.22	32 ± 0.32	
Glycerol	96 ± 0.36	141 ± 0.60	
Detergent	1%	5%	
SDS	7 ± 0.21	6 ± 0.35	
Triton-X-100	86 ± 0.28	83 ± 1.06	
Tween 20 <sup>®</sup>	122 ± 0.36	105 ± 0.52	
Tween 80 <sup>®</sup>	90 ± 0.85	81 ± 0.57	
Inhibitors	5%		
PMSF	27 ± 0.24		
EDTA	30 ± 0.45		
DTT	46 ± 0.30		
β-ME	85 ± 0.56		

which showed that metallic ions can play an important role in maintaining enzymatic structural stability and integrity [50], for example the presence of Fe<sup>3+</sup> and Mn<sup>2+</sup> at 1 mM increased the relative enzymatic activity of the 499EST esterase of *Acidicaldus* USBA-GBX-499 [23]. Moreover, under particular conditions, the presence of metallic cofactors facilitated the correct folding of an enzyme [51] and stabilized the structure of the catalytic site of *Geobacillus zalibae* lipase [52]. The negative effects of metallic ions on the activity of proteins could be explained by undesired interactions occurring between ions and the lateral chains of their amino acids, specifically those involving their surface amino acids, causing ionization of these amino acids [53].




The organic solvents 1-butanol and DMSO inhibited the activity of the 505 LIP. The loss of enzymatic activity in an organic medium has been found to be due to destabilization of the hydrophobic zone of the evaluated protein, and even polar solvents were found to penetrate the protein and induce structural changes [54]. The activity of the enzyme 505 LIP decreased in the presence of acetonitrile, ethanol, 1-propanol and 2-propanol but was not totally inhibited. The lipolytic activity of the 505 LIP was enhanced in the presence of acetone at 15% (v/v), methanol at 15% (v/v) and glycerol at 30% (v/v) (Table 5). The positive effect of these factors on lipolytic activity levels has been reported for other lipolytic enzymes, e.g., the *Pyrococcus furiosus* phospholipase, which has an increased level of enzymatic activity in the presence of methanol and ethanol [55] and the *Aspergillus carneus* thermal-alkaline-resistant esterase, which has an increased level of activity and thermal stability in the presence of glycerol [56]. Stability in the presence of organic solvents is an important property for lipolytic enzymes that are used in non-aqueous systems because it ensures that catalytic reactions can occur under these conditions [57]. This can expand the repertoires of transformations catalysed by lipases, where numerous reactions might be used in the field of synthetic and polymeric chemistry with high selectivity [58]. The presence of organic solvents, such as glycerol in the reaction medium of a lipolytic enzyme, can improve enzymatic activity because they can promote the formation of some hydrogen bonds under anhydrous conditions, generating stable conformations and avoiding structural rigidity and the occurrence of strong electrostatic interactions [59, 60]. Lipases with stability on organic solvents are required for the synthesis of many useful products, such as the production of biopolymeric materials, biodiesel and in the synthesis of fine chemicals [61, 62].

#### *Effect of methanol and glycerol on the stability of the enzyme 505 LIP*

Since methanol at 15 % (v/v) and glycerol at 30 % (v/v) increased the enzymatic activity, the stability of this protein in the presence of those substrates was evaluated (Figure 5). We found that the enzyme 505 LIP not only remained stable in the presence of glycerol for 8 d at room temperature ( $23\text{ }^{\circ}\text{C} \pm 2$ ) but that its activity increased by 50 %. On the other hand, in the presence of methanol, the enzymatic activity increased during the first 24 h of incubation and then decreased by 96 %. The enzymatic stability in the presence of solvents is relevant because they can increase the solubility of non-polar substrates and the thermal stability of enzymes as well as decrease the levels of water-dependent secondary reactions [60, 63]. The polymerization reactions and transesterifications demand harsh conditions provided by the presence of organic solvents and/or high temperature, furthermore industrial lipolytic enzymes are often used in organic media rather than aqueous media [64].

#### *Effect of detergents on the activity of the enzyme 505 LIP*

Regarding the effect of detergents on the enzymatic activity (Table 5), this enzyme exhibited only 10 % of activity in the presence of SDS, in contrast to Triton X-100 and Tween 80, which showed an enzymatic activity of 80 - 90 %. The SDS molecules bind to proteins mainly via hydrophobic interactions, causing the unfolding of their tertiary structures and affecting their activities [65]. In contrast, non-ionic surfactants such as Tween 20<sup>®</sup> increased the enzymatic activity of the 505 LIP by up to 22 %.



**Fig. 5.** Evaluation of lipolytic activity of the enzyme 505 LIP on *p*-nitrophenyl (*p*NP) esters with fatty acid chains of 4 to 18 C-atoms. Activities are shown as percentages of the maximum activity (10.96 U/mg). Values are means  $\pm$  SD from three independent experiments.

Supplementation with a detergent is widely used in biocatalytic settings to resolve pharmaceutical intermediates in non-aqueous media [66]. Additionally, non-ionic surfactants can prevent the denaturation and aggregation of proteins [67]. It has also been reported that Tween 80<sup>®</sup> can adversely affect protein stability by causing oxidative damage generated by residual peroxides of Tween 80<sup>®</sup> that have undergone auto-oxidation [68].

### *Effect of inhibitors on the activity of the enzyme 505 LIP*

The evaluation of the effect of inhibitors on the enzymatic activity (Table 5) shows that EDTA inhibited this activity by 70 % and that PMSF inhibited this activity by 73 %. PMSF is known to inhibit enzymatic activity by affecting the serine in the catalytic site [10], and it is likely that the decrease in the enzymatic activity was caused by modification of an essential serine residue in the active site [63]. The reduced enzymatic activity observed in the presence of the reducing agents DTT and  $\beta$ -ME could be due to the reduction of covalent disulfide bonds between the cysteine residues within this protein, thereby affecting its stability and structure.

## Conclusions

The yeast extract and agitation were key factors in increasing the biomass of *A. facilis* strain USBA-GBX-505 and therefore increasing its growth-associated lipolytic activity. The activity of intracellular lipolytic enzyme from *A. facilis* strain USBA-GBX-505 was

improved by 4.5-fold once the strain was cultured with supplementation of 6.6 g.L<sup>-1</sup> yeast extract and 1 % (v/v) Tween 80<sup>®</sup>, and incubation at 30 °C for 4 d, with agitation at 200 r/min. The partial purification of this enzyme using SEC and IEC provided a recovery yield of 16%. The partially purified enzyme, the 505 LIP, is a moderately thermophilic, intracellular lipase (55 °C) that exhibited the highest enzymatic activity toward middle-chain substrates, particularly *p*NP-C10, at a neutral pH (pH 7.5). The activity of this enzyme is favoured by organic solvents such as glycerol and methanol. This enzyme is stable in the presence of these solvents for a period of 8 d. These characteristics differed from those reported for the esterase produced by *A. facilis* strain AIU409 (called AIU409), suggesting that the enzyme 505 LIP is a novel enzyme produced by a member of the *Acidocella facilis* species. In further studies, the esterification and transesterification reactions of the 505 LIP will be evaluated based on the biochemical characteristics already determined in this study.

## Research and Genetic Resources Access Permits

This study was carried out under the contract MAVDT No. 15, 2008 for genetic resources access (GRA) and the research permit No. DTNO-N-20/2007.

## Conflicts of interest

The authors state that their sole interest in the results of this research is scientific.

## Author contributions

L.F.B and G.L. contributed equally in the results and analysis of this manuscript.

## Acknowledgments

Authors are grateful to Professor Aura Marina Pedroza (Department of Microbiology, Pontificia Universidad Javeriana) for her valuable assistance in the statistical analysis of the data.

## References

- [1] Li S, Yang X, Yang S, Zhu M, Wang X. Technology prospecting on enzymes: application, marketing and engineering, *dsew*, 2: 1-11, 2012.  
doi: [10.5936/csbj.201209017](https://doi.org/10.5936/csbj.201209017)
- [2] Alberty RA. Hydrolase Reactions (Class 3) at 298.15 K, in: *Biochemical Thermodynamics*, John Wiley & Sons, New York, USA, 221-241, 2006.
- [3] Chahinian H, Sarda L. Distinction between esterases and lipases: comparative biochemical properties of sequence-related carboxylesterases, *Protein & Peptide Letters*, 16: 1149-1161, 2009.  
doi: [10.2174/092986609789071333](https://doi.org/10.2174/092986609789071333)
- [4] Sangeetha R, Arulpandi I, Geetha A. Bacterial lipases as potential industrial biocatalysts: an overview, *Research Journal of Microbiology*, 6 :1-24, 2011.-.  
doi: [10.3923/jm.2011.1.24](https://doi.org/10.3923/jm.2011.1.24)

- [5] Godoy Daiha K, Angeli R, Dias de Oliveira S, Volcan Almeida R. Are lipases still important biocatalysts? A study of scientific publications and patents for technological forecasting, *PLoS One*, 10: 6, e0131624, 2015.  
doi: [10.1371/journal.pone.0131624](https://doi.org/10.1371/journal.pone.0131624)
- [6] Joshi S, Satyanarayana T. In vitro engineering of microbial enzymes with multifarious applications: Prospects and perspectives, *Bioresource Technology*, 176: 273-283, 2015.  
doi: [10.1016/j.biortech.2014.10.151](https://doi.org/10.1016/j.biortech.2014.10.151)
- [7] Seghal Kiran G, Nishanth Lipton A, Kennedy J, Dobson ADW, Selvin J. A halotolerant thermostable lipase from the marine bacterium *Oceanobacillus* sp. PUMB02 with an ability to disrupt bacterial biofilms, *Bioengineered*, 5 (5): 305-318, 2014.  
doi: [10.4161/bioe.29898](https://doi.org/10.4161/bioe.29898)
- [8] Masomian M, Rahman RNZRA, Salleh AB, Basri M. A new thermostable and organic solvent-tolerant lipase from *Aneurinibacillus thermoaerophilus* strain HZ, *Process Biochemistry*, 48: 169-175, 2013.  
doi: [10.1016/j.procbio.2012.11.002](https://doi.org/10.1016/j.procbio.2012.11.002)
- [9] Sarkar P, Yamasaki S, Basak S, Bera A, Bag PK, Purification and characterization of a new alkali-thermostable lipase from *Staphylococcus aureus* isolated from *Arachis hypogaea* rhizosphere, *Process Biochemistry*, 47: 858-866, 2012.  
doi: [10.1016/j.procbio.2012.02.023](https://doi.org/10.1016/j.procbio.2012.02.023)
- [10] Kambourova M, Kirilova N, Mandeva R, Dereкова A. Purification and properties of thermostable lipase from a thermophilic *Bacillus stearothermophilus* MC 7, *Journal of Molecular Catalysis B: Enzymatic*, 22: 307-313, 2003.  
doi: [10.1016/S1381-1177\(03\)00045-6](https://doi.org/10.1016/S1381-1177(03)00045-6)
- [11] Sarmiento F, Peralta R, Blamey JM. Cold and hot extremozymes: industrial relevance and current trends, *Frontiers in Bioengineering Biotechnology*, 3: 148, 2015.  
doi: [10.3389/fbioe.2015.00148](https://doi.org/10.3389/fbioe.2015.00148)
- [12] Mazzucotelli CA, Agüero MV, Moreira MdR, Ansorena MR, Optimization of medium components and physicochemical parameters to simultaneously enhance microbial growth and production of lipolytic enzymes by *Stenotrophomonas* sp, *Biotechnology and Applied Biochemistry*.  
doi: [10.1002/bab.1378](https://doi.org/10.1002/bab.1378)
- [13] Swetha S, Varma A, Padmavathi T. Statistical evaluation of the medium components for the production of high biomass,  $\alpha$ -amylase and protease enzymes by *Piriformospora indica* using Plackett–Burman experimental design, *3 Biotech*, 4: 439-445, 2014.  
doi: [10.1007/s13205-013-0168-7](https://doi.org/10.1007/s13205-013-0168-7)
- [14] Gupta R, Gupta N, Rathi P. Bacterial lipases: an overview of production, purification and biochemical properties, *Applied Microbiology Biotechnology*, 64 (6): 763-781, 2004.  
doi: [10.1007/s00253-004-1568-8](https://doi.org/10.1007/s00253-004-1568-8)
- [15] Jones AAB, Philip C. Mineral Microniches Control the Diversity of Subsurface Microbial Populations, *Geomicrobiology Journal*, 31(3) 246-261, 2014.  
doi: [10.1080/01490451.2013.809174](https://doi.org/10.1080/01490451.2013.809174)
- [16] Stierle AA, Stierle DB. Bioprospecting in the Berkeley Pit: The Use of Signal Transduction Enzyme Inhibition Assays to Isolate Bioactive Secondary Metabolites from the Extremophilic Fungi of an Acid Mine Waste Lake, in: Atta-ur-Rahman (Ed.) *Studies in Natural Products Chemistry*, 39: 1-45, 2013.  
doi: [10.1016/B978-0-444-62615-8.00001-1](https://doi.org/10.1016/B978-0-444-62615-8.00001-1)

- [17] Cragg GM, Newman DJ. Natural products: A continuing source of novel drug leads, *Biochimica et Biophysica Acta (BBA) - General Subjects*, 1830 (6): 3670-3695, 2013.  
doi: [10.1016/j.bbagen.2013.02.008](https://doi.org/10.1016/j.bbagen.2013.02.008)
- [18] Zaparty M, Siebers B. Physiology, Metabolism, and Enzymology of Thermoacidophiles, *Extremophiles Handbook*, Springer(Ed.) Japan. pp. 601-639, 2011.  
doi: [10.1007/978-4-431-53898-1\\_28](https://doi.org/10.1007/978-4-431-53898-1_28)
- [19] Amaral-Zettler LA, Zettler ER, Theroux SM, Palacios C, Aguilera A, Amils R. Microbial community structure across the tree of life in the extreme Rio Tinto, *The Isme Journal*, 5: 42-50, 2011.  
doi: [10.1038/ismej.2010.101](https://doi.org/10.1038/ismej.2010.101)
- [20] G. López, Composición de la diversidad microbiana y búsqueda de enzimas lipolíticas en ambientes ácidos del parque nacional natural de Los Nevados, Tesis Doctoral, Facultad de Ciencias. Departamento de Biología, Pontificia Universidad Javeriana, pp. 280, 2013.
- [21] Itoh T, Suzuki K, Sanchez PC, Nakase T. *Caldisphaera lagunensis* gen. nov., sp. nov., a novel thermoacidophilic crenarchaeote isolated from a hot spring at Mt Maquiling, Philippines, *International Journal of Systematic and Evolutionary Microbiology*, 53: 1149-1154, 2003.  
doi: [10.1099/ijs.0.02580-0](https://doi.org/10.1099/ijs.0.02580-0)
- [22] Imhoff-Stuckle D, Pfennig N, Isolation and characterization of a nicotinic acid-degrading sulfate-reducing bacterium, *Desulfococcus niacini* sp. nov., *Archives of Microbiology*, 136: 194-198, 1983.
- [23] Lopez G, Chow J, Bongen P, Lauinger B, Pietruszka J, Streit WR, Baena S. A novel thermoalkalostable esterase from *Acidicaldus* sp. strain USBA-GBX-499 with enantioselectivity isolated from an acidic hot springs of Colombian Andes, *Applied Microbiology Biotechnology*, 98: 8603-8616, 2014.  
doi: [10.1007/s00253-014-5775-7](https://doi.org/10.1007/s00253-014-5775-7)
- [24] Bradford M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding, *Analytical Biochemistry*, 72: 248-254, 1976.  
doi: [10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
- [25] Rajendran A, Thangavelu V. Application of Central Composite Design and Artificial Neural Network for the Optimization of Fermentation Conditions for Lipase Production by *Rhizopus arrhizus* MTCC 2233, *Journal of Bioprocessing & Biotechniques*, 2:118  
doi: [10.4172/2155-9821.1000118](https://doi.org/10.4172/2155-9821.1000118)
- [26] Arevalos-Sánchez M, Regalado C, Martin SE, Domínguez-Domínguez J, García-Almendárez BE. Effect of neutral electrolyzed water and nisin on *Listeria monocytogenes* biofilms, and on listeriolysin O activity, *Food Control*, 24: 116-122, 2012.  
doi: [10.1016/j.foodcont.2011.09.012](https://doi.org/10.1016/j.foodcont.2011.09.012)
- [27] Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4, *Nature*, 227: 680-685, 1970.  
doi: [10.1038/227680a0](https://doi.org/10.1038/227680a0)
- [28] Switzer II RC, Merrill CR, Shifrin S. A highly sensitive silver stain for detecting proteins and peptides in polyacrylamide gels, *Analytical Biochemistry*, 98: 231-237, 1979.  
doi: [10.1016/0003-2697\(79\)90732-2](https://doi.org/10.1016/0003-2697(79)90732-2)
- [29] Niyonzima FN, More SS. Concomitant production of detergent compatible enzymes by *Bacillus flexus* XJU-1, *Brazilian Journal of Microbiology*, 45(3): 903-910,2014.  
Retrieved from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4204975/>

- [30] Sharma CK, Sharma PK, Kanwar SS. Optimization of production conditions of lipase from *B. licheniformis* MTCC-10498, *Research Journal of Recent Sciences*, 1(7): 23-32, 2012.  
Retrieved from:  
<http://www.isca.in/rjrs/archive/v1/i7/4.ISCA-RJRS-2012-140%20Done.php>
- [31] Abol Fotouh DM, Bayoumi RA, Hassan MA. Production of Thermoalkaliphilic Lipase from *Geobacillus thermoleovorans* DA2 and Application in Leather Industry, *Enzyme Research*, 2016 (2016) 9034364.  
doi: [10.1155/2016/9034364](https://doi.org/10.1155/2016/9034364)
- [32] K. Isobe, N. Wakao, Purification and some properties of a thermostable acid esterase from *Acidiphilium* sp. AIU 409, *The Journal of General Applied Microbiology*, 43 (3): 151-156, 1997.  
doi: [doi.org/10.2323/jgam.43.151](https://doi.org/10.2323/jgam.43.151)
- [33] Kishimoto N, Kosako Y, Wakao N, Tano T, Hiraishi A. Transfer of *Acidiphilium facilis* and *Acidiphilium aminolytica* to the Genus *Acidocella* gen. nov., and Emendation of the Genus *Acidiphilium*, *Systematic and Applied Microbiology*, 18 (1):85-91, 1995.  
doi: [10.1016/S0723-2020\(11\)80453-4](https://doi.org/10.1016/S0723-2020(11)80453-4)
- [34] Barrie Johnson D, Hallberg KB. Carbon, iron and sulfur metabolism in acidophilic microorganisms, *Advance in Microbial Physiology*, 54: 201-255, 2009.  
doi: [10.1016/S0065-2911\(08\)00003-9](https://doi.org/10.1016/S0065-2911(08)00003-9)
- [35] Tanguler H, Erten H. Utilisation of spent brewer's yeast for yeast extract production by autolysis: the effect of temperature, *Food and Bioproducts Processing*, 86: 317-321, 2008.  
doi: [10.1016/j.fbp.2007.10.015](https://doi.org/10.1016/j.fbp.2007.10.015)
- [36] Salihu A, Alam MZ, AbdulKarim MI, Salleh HM, Optimization of lipase production by *Candida cylindracea* in palm oil mill effluent based medium using statistical experimental design, *Journal of Molecular Catalysis B: Enzymatic*, 69: 66-73, 2011.  
doi: [10.1016/j.molcatb.2010.12.012](https://doi.org/10.1016/j.molcatb.2010.12.012)
- [37] Liu CH, Lu WB, Chang JS. Optimizing lipase production of *Burkholderia* sp. by response surface methodology, *Process Biochemistry*, 41: 1940-1944, 2006.  
doi: [10.1016/j.procbio.2006.04.013](https://doi.org/10.1016/j.procbio.2006.04.013)
- [38] Liu CH, Chen CY, Wang YW, Chang JS. Fermentation strategies for the production of lipase by an indigenous isolate *Burkholderia* sp. C20, *Biochemical Engineering Journal*, 58-59: 96-102, 2011.  
doi: [10.1016/j.bej.2011.09.001](https://doi.org/10.1016/j.bej.2011.09.001)
- [39] Chen JY, Wen CM, Chen TL. Effect of oxygen transfer on lipase production by *Acinetobacter radioresistens*, *Biotechnology and Bioengineering*, 62: 311-316, 1999.
- [40] Boekema BKHL, Beselin A, Breuer M, Hauer B, Koster M, Rosenau F, Jaeger KE, Tommassen J. Hexadecane and Tween 80 stimulate lipase production in *Burkholderia glumae* by different mechanisms, *Applied Environmental Microbiology*, 73: 3838-3844, 2007.  
doi: [10.1128/AEM.00097-07](https://doi.org/10.1128/AEM.00097-07)
- [41] Kamarudin NH, Rahman RN, Ali MS, Leow TC, Basri M, Salleh AB. A new cold-adapted, organic solvent stable lipase from mesophilic *Staphylococcus epidermidis* AT2, *The Protein Journal*, 33 296-307, 2014.  
doi: [10.1007/s10930-014-9560-3](https://doi.org/10.1007/s10930-014-9560-3)
- [42] Li CY, Cheng CY, Chen TL. Production of *Acinetobacter radioresistens* lipase using Tween 80 as the carbon source, *Enzyme and Microbial Technology*, 29: 258-263, 2001.  
doi: [10.1016/S0141-0229\(01\)00396-9](https://doi.org/10.1016/S0141-0229(01)00396-9)

- [43] Saranya P, Sukanya Kumari H, Prasad Rao B, Sekaran G. Lipase production from a novel thermo-tolerant and extreme acidophile *Bacillus pumilus* using palm oil as the substrate and treatment of palm oil-containing wastewater, *Environmental Science and Pollution Research*, 21: 3907-3919, 2014.  
doi: [10.1007/s11356-013-2354-x](https://doi.org/10.1007/s11356-013-2354-x)
- [44] Contesini FJ, Lopes DB, Macedo GA, Nascimento MdG, Carvalho PdO. *Aspergillus* sp. lipase: potential biocatalyst for industrial use, *Journal of Molecular Catalysis B: Enzymatic*, 67: 163-171, 2010.  
doi: [10.1016/j.molcatb.2010.07.021](https://doi.org/10.1016/j.molcatb.2010.07.021)
- [45] Takahashi T, Tanaka N, Tamura T, Mukai C, Isobe K, Wakao N, Tanaka H, Inagaki K, Cloning of the extracellular acid esterase gene from acidophilic bacterium, *Acidocella facilis*, *Scientific Reports of the Faculty of Agriculture*, Okayama University, 91: 7-13, 2002.  
Retrieved of <http://ousar.lib.okayama-u.ac.jp/metadata/10603>
- [46] Lesuisse E, Schanck K, Colson C. Purification and preliminary characterization of the extracellular lipase of *Bacillus subtilis* 168, an extremely basic pH-tolerant enzyme, *European Journal Biochemistry*, 216: 155-160, 1993.  
doi: [10.1111/j.1432-1033.1993.tb18127.x](https://doi.org/10.1111/j.1432-1033.1993.tb18127.x)
- [47] Tamilarasan K, Dharmendira Kumar M. Purification and characterization of solvent tolerant lipase from *Bacillus sphaericus* MTCC 7542, *Biocatalysis and Agricultural Biotechnology*, 309-313, 2012.  
doi: [10.1016/j.bcab.2012.07.001](https://doi.org/10.1016/j.bcab.2012.07.001)
- [48] Mandrich L, Menchise V, Alterio V, De Simone G, Pedone C, Rossi M, Manco G. Functional and structural features of the oxyanion hole in a thermophilic esterase from *Alicyclobacillus acidocaldarius*, *Proteins*, 71: 1721-1731, 2008.  
doi: [10.1002/prot.21877](https://doi.org/10.1002/prot.21877)
- [49] Auernik KS, Cooper CR, Kelly RM. Life in hot acid: pathway analyses in extremely thermoacidophilic archaea, *Current Opinion in Biotechnology*, 19: 445-453, 2008.  
doi: [10.1016/j.copbio.2008.08.001](https://doi.org/10.1016/j.copbio.2008.08.001)
- [50] Çolak A, Sisik D, Sağlam N, Güner S, Çanakçı S, Beldüz AO. Characterization of a thermoalkalophilic esterase from a novel thermophilic bacterium, *Anoxybacillus gonensis* G2, *Bioresource Technology*, 96: 625-631, 2005.  
doi: [10.1016/j.biortech.2004.06.003](https://doi.org/10.1016/j.biortech.2004.06.003)
- [51] Bushmarina NA, Blanchet CE, Vernier G, Forge V. Cofactor effects on the protein folding reaction: acceleration of  $\alpha$ -lactalbumin refolding by metal ions, *Protein Science*, 15: 659-671, 2006.  
doi: [10.1110/ps.051904206](https://doi.org/10.1110/ps.051904206)
- [52] Wahab R.A, Basri M, Rahman MBA, Rahman RNZRA, Salleh A.B., Chor L.T., Engineering catalytic efficiency of thermophilic lipase from *Geobacillus zalihae* by hydrophobic residue mutation near the catalytic pocket, *Advances in Bioscience and Biotechnology*, 3: 158-167, 2012.  
doi: [10.4236/abb.2012.32024](https://doi.org/10.4236/abb.2012.32024)
- [53] Dandavate V, Jinjala J, Keharia H, Madamwar D, Production, partial purification and characterization of organic solvent tolerant lipase from *Burkholderia multivorans* V2 and its application for ester synthesis, *Bioresource Technology*, 100: 3374-3381, 2009  
doi: [10.1016/j.biortech.2009.02.011](https://doi.org/10.1016/j.biortech.2009.02.011)
- [54] Maytorena-Verdugo C, García-Carreño F. Uso de enzimas en solventes orgánicos en la industria de bioprocesos, *BioTecnología*, 16: 15, 2012

- [55] Chandrayan SK, Dhaunta N, Guptasarma P. Expression, purification, refolding and characterization of a putative lysophospholipase from *Pyrococcus furiosus*: retention of structure and lipase/esterase activity in the presence of water-miscible organic solvents at high temperatures, *Protein Expression and Purification*, 59: 327-333, 2008.  
doi: [10.1016/j.pep.2008.02.019](https://doi.org/10.1016/j.pep.2008.02.019)
- [56] Saxena RK, Davidson WS, Sheoran A, Giri B. Purification and characterization of an alkaline thermostable lipase from *Aspergillus carneus*, *Process Biochemistry*, 39: 239-247, 2003.  
doi: [10.1016/S0032-9592\(03\)00068-2](https://doi.org/10.1016/S0032-9592(03)00068-2)
- [57] Doukyu N, Ogino H. Organic solvent-tolerant enzymes, *Biochemistry Engineering Journal*, 48: 270-282, 2010.  
doi: [10.1016/j.bej.2009.09.009](https://doi.org/10.1016/j.bej.2009.09.009)
- [58] Souza LT, Oliveira JS, dos Santos VL, Regis WC, Santoro MM, Resende RR. Lipolytic potential of *Aspergillus japonicus* LAB01: production, partial purification, and characterisation of an extracellular lipase, *Biomed Research International*, 2014: 108913, 2014.  
doi: [10.1155/2014/108913](https://doi.org/10.1155/2014/108913)
- [59] Faulds CB, Pérez-Boada M, Martínez ÁT. Influence of organic co-solvents on the activity and substrate specificity of feruloyl esterases, *Bioresource Technology*, 102: 4962-4967, 2011.  
doi: [10.1016/j.biortech.2011.01.088](https://doi.org/10.1016/j.biortech.2011.01.088)
- [60] Herbst D, Peper S, Niemeyer B. Enzyme catalysis in organic solvents: influence of water content, solvent composition and temperature on *Candida rugosa* lipase catalyzed transesterification, *Journal of Biotechnology*, 162: 398-403, 2012.  
doi: [10.1016/j.jbiotec.2012.03.011](https://doi.org/10.1016/j.jbiotec.2012.03.011)
- [61] Dahiya P, Chand S, Dilbaghi N. Immobilization of Organic Solvent-Tolerant Lipase from *Pseudomonas mendocina* M-37 with Potential Synthetic Activities, *Food Technology Biotechnology*, 52(3): 368-375, 2014.  
Retrieved from:  
[http://www.ftb.com.hr/images/pdfarticles/2014/July-September/ftb\\_52-3\\_368-375.pdf](http://www.ftb.com.hr/images/pdfarticles/2014/July-September/ftb_52-3_368-375.pdf)
- [62] Pellis A, Herrero Acero E, Gardossi L, Ferrario V, Guebitz GM. Renewable building blocks for sustainable polyesters: new biotechnological routes for greener plastics, *Polymer International*, 65 (8): 861-871, 2016.  
doi: [10.1002/pi.5087](https://doi.org/10.1002/pi.5087)
- [63] Kumar L, Singh B, Adhikari DK, Mukherjee J, Ghosh D. A thermoalkaliphilic halotolerant esterase from *Rhodococcus* sp. LKE-028 (MTCC 5562): enzyme purification and characterization, *Process Biochemistry*, 47: 983-991, 2012.  
doi: [10.1016/j.procbio.2012.03.020](https://doi.org/10.1016/j.procbio.2012.03.020)
- [64] Sharma S, Kanwar SS. Organic solvent tolerant lipases and applications, *The Scientific World Journal*, 2014: 625258, 2014.  
doi: [10.1155/2014/625258](https://doi.org/10.1155/2014/625258)
- [65] Bhuyan A.K. On the mechanism of SDS-induced protein denaturation, *Biopolymers*, 93: 186-199, 2010.  
doi: [10.1002/bip.21318](https://doi.org/10.1002/bip.21318)
- [66] Cao Y, Zhuang Y, Yao C, Wu B, He B. Purification and characterization of an organic solvent-stable lipase from *Pseudomonas stutzeri* LC2-8 and its application for efficient resolution of (R, S)-1-phenylethanol, *Biochemical Engineering Journal*, 64: 55-60, 2012.  
doi: [10.1016/j.bej.2012.03.004](https://doi.org/10.1016/j.bej.2012.03.004)



- [67] Chou DK, Krishnamurthy R, Randolph TW, Carpenter JF, Manning MC. Effects of Tween 20<sup>®</sup> and Tween 80<sup>®</sup> on the stability of Albutropin during agitation, *Journal Pharmaceutical Sciences*, 94: 1368-1381, 2005.  
doi: [10.1002/jps.20365](https://doi.org/10.1002/jps.20365)
- [68] Wang W, Wang YJ, Wang D. Dual effects of Tween 80 on protein stability, *International Journal Pharmaceutics*, 347: 31-38, 2008.  
doi: [10.1016/j.ijpharm.2007.06.042](https://doi.org/10.1016/j.ijpharm.2007.06.042)

### Metodología de Superficie de Respuesta (RSM) para el análisis de las condiciones de cultivo de *Acidocella facilis* cepa USBA-GBX-505 y la purificación parcial y caracterización bioquímica de la lipasa 505 LIP

**Resumen.** Por medio de la Metodología de Respuesta de Superficie (RSM) evaluamos las condiciones de cultivo (fuente de N, fuente de C, pH y tasa de agitación) que incrementan la biomasa de *Acidocella facilis* cepa USBA-GBX-505 y, como consecuencia, la producción de su enzima lipolítica, llamada 505 LIP. Los resultados de la RSM revelaron que el extracto de levadura y la agitación fueron factores de cultivo claves, que incrementaron de 4 a 5 veces la actividad lipolítica asociada al crecimiento (de 0.13 U.mg<sup>-1</sup> a 0.6 U.mg<sup>-1</sup>). La lipasa 505 LIP se purificó parcialmente usando cromatografía de exclusión por tamaño y cromatografía de intercambio iónico. Su peso molecular fue > 77 kDa. La enzima muestra su actividad catalítica óptima a 55 °C y pH 7.5. El EDTA, el PMSF, el 1-butanol y el DMSO inhibieron la actividad enzimática, mientras que el Tween 20, la acetona, el glicerol y el metanol la incrementaron. La enzima 505 LIP no requiere iones metálicos para su actividad, e incluso se inhibe en presencia de ellos. Este estudio muestra el uso potencial de *A. facilis* cepa USBA-GBX-505 para la producción de una nueva enzima lipolítica, 505 LIP, que es estable a temperaturas moderadas y en la presencia de solventes orgánicos. Estas son características importantes en la síntesis de muchos productos útiles.

**Palabras clave:** lipasas; acidófilos; *Acidocella facilis*; estabilidad en solventes; diseño Plackett-Burman; diseño central compuesto.

### Metodologia de Superficie de Resposta (MSR) para a análise das condições de cultivo de *Acidocella facilis* cepa USBA-GBX-505 e purificação parcial e caracterização bioquímica da lipase 505 LIP.

**Resumo.** Utilizando a Metodologia de Superfície de Resposta (MSR) avaliamos as condições de cultivo (fontes de nitrogênio e carbono, pH e taxa de agitação) que aumentam a biomassa de *Acidocella facilis* cepa USBA-GBX-505, e, portanto, elevam a produção de sua enzima lipolítica 505 LIP. Os resultados da MSR revelaram que o extrato de levedura e a agitação foram fatores de cultivo chave que permitiram aumentar 4 a 5 vezes a atividade lipolítica associada ao crescimento (de 0,13 U.mg<sup>-1</sup> a 0,6 U.mg<sup>-1</sup>). A lipase 505 LIP foi parcialmente purificada utilizando cromatografia por exclusão de tamanho e cromatografia de intercâmbio iônico. Seu peso molecular foi > 77 kDa. A enzima mostra sua atividade catalítica ótima a 55 °C e pH 7,5. EDTA, PMSF, 1-butanol e DMSO inibiram a atividade enzimática, enquanto que Tween 20, acetona, glicerol e metanol aumentaram esta atividade. Íons metálicos não são necessários para a atividade da 505 LIP, apresentando inclusive efeito inibitório da enzima. Este estudo demonstra o potencial uso de *A. facilis* cepa USBA-GBX-505 para a produção de uma nova enzima lipolítica, 505 LIP, a qual é estável a moderadas temperaturas e na presença de solventes orgânicos. Estas características são importantes para a síntese de diversos produtos úteis.

**Palavras-chave:** lipases; acidófilos; *Acidocella facilis*; estabilidade em solventes; experimento Plackett-Burman; experimento central composto.



#### Luisa Fernanda Bernal, M.Sc.

She has a Bachelor degree on Industrial microbiology and a Master on Science at the Pontificia Universidad Javeriana. Currently, she is a professor at Universidad Nacional Abierta y a Distancia. She also is a consultant in Quality Management Systems especially NTC/ISO 17025. Since 2015 she is studying laws at Universidad del Sinú. Her professional interests focus on biotechnology, intellectual property and environmental laws.



#### Gina López Ramírez, Ph.D.

She is an industrial Microbiologist and a PhD in Sciences at Pontificia Universidad Javeriana. Currently, she is a postdoctoral researcher in the Department of Biology at the Pontificia Universidad Javeriana. She has been working on microbial diversity of extremophilic microorganisms (acidophiles, thermophiles, psychrophiles), focused on lipolytic enzymes from these microorganisms. Other research areas are development of transformation reactions of conifer oleoresins, triacylglycerols and fatty acids using lipolytic enzymes for generating intermediate substrates. Actually she is working on characterization of lipids from acidophilic microalgae.



#### Monica Irene Ruiz Camacho, Msc.

Monica has a bachelor degree on Chemistry from the Universidad Industrial de Santander with a Master's degree in Chemistry at the same university. Actually she is a research assistant in the Department of Biology at Pontificia Universidad Javeriana. The research areas are biotechnology, bio-processes and enzymatic catalysts.

#### Alejandro Reyes Cardenas

Chemical Engineer at Universidad de America, he worked at the Chemistry laboratory of the Attorney General of the Nation during 14 years and currently he works at the Chemistry Department in the chromatography laboratories of the Pontificia Universidad Javeriana.

#### Ricardo Vera Bravo, Ph.D.

Is a chemist and received Ph.D. in chemistry from Universidad Nacional de Colombia. He worked with Fundación Instituto de Inmunología de Colombia (12 years) and is currently professor of the Chemistry Department at Pontificia Universidad Javeriana, where is teaching analytical chemistry and organic chemistry. His professional interests focus on isolation, purification and the determination of biological activity of macromolecules from plant and animal extracts. The research projects include isolation of GMP as control of adulteration from milk, purification of lipolytic enzymes, peptides with antimicrobial activity, determination of cytotoxic activity in tropical fruits.



#### Sandra Baena Garzón, Msc, Ph.D.

Associate Professor in the Department of Biology at Pontificia Universidad Javeriana. She is a biologist at the Pontificia Universidad Javeriana, she has a Master's degree in Sanitation and Environmental Development at the same university and a PhD in Sciences from the University of Provence, Aix-Marseille I (France). The research areas are diversity and microbial ecology of anaerobic wastewater treatment and thermal and saline springs. She was Member of the National Biotechnology Council in COLCIENCIAS and she is Corresponding member of the Academia Colombiana de Ciencias Exactas, Físicas y Naturales.