

## Simple preparation of new *N*-(6-methyl-2-nitrophenyl-1,2,3,4-tetrahydroquinolin-4-yl) pyrrolidin-2-ones and their spectroscopic analysis

John H. Bermúdez<sup>1</sup>, Jose Luis Pinto<sup>2</sup>, Carlos Mario Meléndez<sup>3</sup>, José A. Henao<sup>2\*</sup>, Vladímir V. Kouznetsov<sup>1\*</sup>

<sup>1</sup>Laboratorio de Química Orgánica y Biomolecular, Escuela de Química, Universidad Industrial de Santander, A.A. 678, Bucaramanga, Colombia.

<sup>2</sup>Grupo de Investigación en Química Estructural, CIBIMOL, Escuela de Química, Universidad Industrial de Santander, A.A. 678, Bucaramanga, Colombia.

<sup>3</sup>Grupo de Investigación en Compuestos Heterocíclicos, Programa de Química, Facultad de Ciencias Básicas, Universidad del Atlántico, A.A.1890, Barranquilla, Colombia.

\* kouznet@uis.edu.co ; jahenao@uis.edu.co

Received: 16-05-2011; Accepted: 05-07-2011

### Abstract

**Objectives.** To prepare new *N*-(1,2,3,4-tetrahydroquinolin-4-yl) pyrrolidin-2-one molecules and to characterize them by spectroscopic methods. **Materials and methods.** All reagents were purchased from Aldrich, commercial grade. The purity of the products and the composition of the reaction mixtures were monitored by thin layer chromatography over Silufol UV<sub>254</sub> chromatoplates (0.25 mm). Product isolation and purification were performed by column chromatography (SiO<sub>2</sub>) using ethyl acetate. **Results.** Preparation of new *N*-(2-nitrophenyl-1,2,3,4-tetrahydroquinolin-4-yl) pyrrolidin-2-ones has been achieved via the one-pot synthesis, based on a BiCl<sub>3</sub>-catalyzed imino Diels-Alder cycloaddition reaction of toluidine, *N*-vinylpyrrolidin-2-one and 4-nitro- or 3-nitrobenzaldehydes. The structure of the pyrrolidine derivatives was confirmed by <sup>1</sup>H NMR and <sup>13</sup>C NMR studies, in addition to inverse-detected 2D NMR experiments and monocrystal X-ray diffraction. **Conclusions.** An efficient, economic, and fast synthetic route (multi-component imino Diels-Alder reaction) was employed in the construction of several new tetrahydroquinoline derivatives, useful and attractive rigid skeleton with well-defined stereochemistry.

**Key words:** tetrahydroquinoline derivatives, *N*-substituted pyrrolidin-2-ones, three component imino Diels-Alder reaction, one-pot synthesis.

### Resumen

**Preparación simple de nuevas *N*-(6-metil-2-nitrofenil-1,2,3,4-tetrahydroquinolin-4-il) pirrolidin-2-onas y su análisis espectroscópico.**

**Objetivos.** Preparar nuevas moléculas *N*-(1,2,3,4-tetrahydroquinolin-4-il) 2-oxipirrolidínicas y caracterizarlas por métodos espectroscópicos.

**Materiales y métodos.** Todos los reactivos usados son de Aldrich, grado comercial. La pureza de los productos y la composición de las mezclas de reacción fueron monitoreadas por cromatografía en capa fina sobre cromatoplasmas de Silufol UV<sub>254</sub> (0.25 mm). El aislamiento y purificación se realizó usando cromatografía en columna (SiO<sub>2</sub>), usando acetato de etilo. **Resultados.** La preparación de las nuevas *N*-(tetrahydroquinolin-4-il) pirrolidin-2-onas 4-nitrofenil (ó 2-nitrofenil) sustituidas en C-2 del anillo tetrahydroquinolínico, se realizó vía síntesis *one-pot* basada en la reacción de cicloadición imino Diels-Alder catalizada por BiCl<sub>3</sub> entre toluidina, *N*-vinilpirrolidin-2-ona y 4-nitrobenzaldehído (3-nitrobenzaldehído). La estructura de los derivados pirrolidínicos fue confirmada por <sup>1</sup>H RMN y <sup>13</sup>C RMN, además de experimentos 2D RMN y difracción de rayos X de monocristal. **Conclusiones.** Una ruta eficiente, económica y rápida (reacción imino Diels-Alder multi-componente) fue empleada para la construcción de nuevas moléculas *N*-(tetrahydroquinolin-4-il) 2-oxipirrolidínicas, esqueleto muy atractivo y usado con estereoquímica bien definida.

**Palabras clave:** derivados de la 1,2,3,4-tetrahydroquinolina, pirrolidin-2-onas *N*-sustituidas, reacción imino Diels-Alder de tres componentes, síntesis *one-pot*.

## Resumo

**Preparação simples de novas N-(6-metil-2-nitrofenil-1,2,3,4-tetrahydroquinoline-4-il) pirrolidin-2-onas e sua análise espectroscópica.**

**Objetivos.** Preparar novas moléculas N-(1,2,3,4-tetrahydroquinoline-4-il) 2-oxopirrolidínicas e sua caracterização por espectroscopia.

**Materiais e métodos.** Todos os reagentes utilizados são de Aldrich, de grau comercial. A pureza dos produtos e a composição das misturas de reação foram monitoradas por cromatografia em camada fina sobre cromatoplasmas de Silufol UV<sub>254</sub> (0,25 mm). O isolamento e purificação foi realizado utilizando cromatografia em coluna (SiO<sub>2</sub>), utilizando acetato de etila. **Resultados.** Preparação de novas N-(tetrahydroquinoline-4-il) pirrolidin-2-onas 4-nitrofenil (ou 2-nitrofenil) substituídas em C-2 do anel tetrahydroquinoline foi realizada através da síntese “one pot” baseada na reação de cicloadição imino Diels-Alder catalisada por BiCl<sub>3</sub> entre toluidina, N-vinilpirrolidin-2-ona e 4 nitrobenzaldehyde (3 nitrobenzaldehyde). A estrutura dos derivados pirrolidínicos foi confirmada por <sup>1</sup>H RMN y <sup>13</sup>C RMN, experimentos 2D RMN, assim como difração de raios X e monocristais. **Conclusões.** Uma rota eficiente, econômica e rápida (reação imino Diels-Alder multi-componente) foi utilizada para a construção de novas moléculas N-(tetrahydroquinoline-4-il) 2-oxopirrolidínicas esqueleto muito atraente e usado com estereoquímica bem definida.

**Palavras-chave:** derivados de 1,2,3,4-tetrahydroquinoline, pirrolidin-2-onas N-substituídas, reação imino Diels-Alder de três componentes, síntese “one-pot”.

## Introduction

Quinoline and tetrahydroquinoline structures are an essential feature of many natural products. These heterocycles play a key role in heterocyclic and medicinal chemistry. Their synthesis by various methodologies has been published extensively (1-4). Polyfunctionalized tetrahydroquinolines (THQs) are molecules of great interest in organic synthesis since many natural products present this system in their structure, and because they exhibit diverse biological activities (5-9). Besides their notorious bioactivity, THQs are also important and reliable precursors in quinoline preparation, another group of heterocyclic molecules that has a great number of pharmacological properties (10). An efficient route in the preparation of THQs is the acid-catalyzed Povarov reaction that is classified as imino Diels-Alder cycloaddition (11-13) and permits the condensation of anilines, aldehydes, and electron-rich alkenes using acidic catalysts under mild conditions to achieve the obtainment of new substituted tetrahydroquinolines.

As a part of our research program in the DOS methodology towards the synthesis of bioactive substituted tetrahydroquinolines and quinolines, we are currently conducting research on the synthesis of small drug-like (tetrahydro)quinoline molecules containing a C-2 aryl fragment, whose synthesis could be accomplished via cycloaddition reactions. We want to report here the simple preparation of new N-(2-nitrophenyl-1,2,3,4-tetrahydroquinolin-4-yl) pyrrolidin-2-ones using BiCl<sub>3</sub>-catalyzed three-component Povarov reaction among nitrobenzaldehydes, toluidine and N-vinylpyrrolidin-2-one, and their transformations into potentially bioactive 2-aryl-tetrahydroquinoline derivatives, N-amidyl substituted at the C-4 position.

## Materials and methods

All reagents were purchased from Aldrich, commercial grade. The purity of the products and the composition of the reaction mixtures were monitored by thin layer chromatography over Silufol UV<sub>254</sub> 0.25 mm-thick chromatoplates. The melting points (uncorrected) were determined on a Fisher-Johns melting point apparatus. The IR spectra were recorded on an Infracum FT-02 spectrophotometer in KBr. <sup>1</sup>H NMR spectra were recorded on Bruker AM-400 or AC-300 spectrometers in CDCl<sub>3</sub>. Chemical shifts are reported in ppm. (A signal at 7.24 ppm of CHCl<sub>3</sub> in CDCl<sub>3</sub> was used as reference for protons). A Hewlett Packard 5890a series II Gas Chromatograph interfaced to an HP 5972 Mass Selective Detector (MSD) with an HPMS Chemstation Data System was used for MS identification at 70 eV using a 60 m capillary column coated with HP-5 [5%-phenyl-poly(dimethyl-siloxane)]. X ray diffraction single-crystal technique with an AFC7S four circle diffractometer was used. The data acquisition was made to 293 K of temperature with MoKa (λ = 0.71073 Å) radiation and a measurement range between 1 and 25° of theta (θ). The structure elucidation and the refinement were made with the software Shelxs-97 and Shelxl-97, respectively. Elemental analyses were performed on a Perkin Elmer 2400 Series II analyzer and were within ± 0.4 of the theoretical values. The reaction progress was monitored using thin layer chromatography on a silufol UV<sub>254</sub> TLC aluminum sheet.

## Synthesis of new N-(2-nitrophenyl-1,2,3,4-tetrahydroquinolin-4-yl) pyrrolidin-2-ones. General procedure

20 mol% BiCl<sub>3</sub> was added to a solution of the toluidine 1 (2.85 mmol) and an appropriate nitrobenzaldehyde **3** or **4** (3.13 mmol) in anhydrous CH<sub>3</sub>CN (15 mL) under N<sub>2</sub>, and

*N*-vinylpyrrolidin-2-one **2** (3.42mmol) was added to the resulting mixture. The reaction mixture was stirred at room temperature for 20-24 h and then quenched with a solution of Na<sub>2</sub>CO<sub>3</sub>. The organic layer was separated and dried with Na<sub>2</sub>SO<sub>4</sub>. The organic solvent was removed in vacuum to obtain the respective *N*-(2-nitrophenyl-1,2,3,4-tetrahydroquinolin-4-yl) pyrrolidin-2-ones **5** or **6**. The reaction mixture was adsorbed under silica gel and separated by chromatography (Hexane / Ethyl acetate).

### **N**-[6-Methyl-2-(4-nitrophenyl)-1,2,3,4-tetrahydroquinolin-4-yl] pyrrolidin-2-one (**5**)

Yellow solid. Mp. 222-223°C. Yield 95%. Anal. calcd for C<sub>20</sub>H<sub>21</sub>N<sub>3</sub>O<sub>3</sub>: C, 68.35; H, 6.00; N, 11.99. M = 351.40. Found: C, 68.46; H, 6.22; N, 12.06. GC-MS: R<sub>t</sub> = 44.57 min; *m/z* (EI): 351 (M<sup>+</sup>). IR (KBr): ν 3394, 2916, 1666 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.20 (2H, d, *J* = 8.7 Hz, 3'-H and 5'-H), 7.61 (2H, d, *J* = 8.7 Hz, 2'-H and 6'-H), 6.90 (1H, dd, *J* = 8.0, 1.7 Hz, 7-H), 6.68 (1H, s, 5-H), 6.57 (1H, d, *J* = 8.1 Hz, 8-H), 5.69 (1H, dd, *J* = 11.1, 6.4 Hz, 4-H<sub>ax</sub>), 4.65 (1H, dd, *J* = 10.7, 3.1 Hz, 2-H<sub>ax</sub>), 4.03 (1H, br.s, NH), 3.21 (2H, t, *J* = 6.9 Hz, 5''-H), 2.59-2.41 (2H, m, 3''-H), 2.23 (3H, s, 6-CH<sub>3</sub>), 2.13-1.99 (4H, m, 4''-H and 3-H) ppm. <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 175.8, 150.6, 147.4, 142.9, 129.0, 128.1, 127.3 (2C), 126.9, 123.9 (2C), 118.8, 115.4, 56.0, 48.1, 42.2, 35.3, 31.3, 20.5, 18.1 ppm.

### **N**-[6-Methyl-2-(3-nitrophenyl)-1,2,3,4-tetrahydroquinolin-4-yl] pyrrolidin-2-one (**6**)

Yellow solid. Mp. 242-243 °C. Yield 70%. Anal. calcd for C<sub>20</sub>H<sub>21</sub>N<sub>3</sub>O<sub>3</sub>: C, 68.36; H, 6.02; N, 11.96. M = 351.40. Found: C, 68.57; H, 6.28; N, 11.74. GC-MS: R<sub>t</sub> = 45.32 min; *m/z* (EI) 264 (M<sup>+</sup>-87). IR (KBr): ν 3326, 2900, 1630 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.41 (1H, t, *J* = 2.0 Hz, 2'-H), 8.20 (1H, ddd, *J* = 8.1, 2.3, 1.0 Hz, 6'-H), 7.77 (1H, br.d, *J* = 7.8 Hz, 4'-H), 7.57 (1H, t, *J* = 7.8 Hz, 5'-H), 6.93 (1H, dd, *J* = 8.1, 2.0 Hz, 7-H), 6.72 (1H, s, 5-H), 6.60 (1H, d, *J* = 8.1 Hz, 8-H), 5.74 (1H, dd, *J* = 11.2, 7.1 Hz, 4-H), 4.71 (1H, dd, *J* = 10.4, 3.5 Hz, 2-H), 4.01 (1H, br.s, NH), 3.30-3.20 (2H, m, 5''-H<sub>pyrr</sub>), 2.62-2.45 (2H, m, 3''-H<sub>pyrr</sub>), 2.26 (3H, s, 6-CH<sub>3</sub>), 2.16-2.13 (2H, m, 3-H), 2.10-2.02 (2H, m, 4''-H<sub>pyrr</sub>) ppm. <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 175.9, 148.7, 145.4, 132.9, 129.9, 128.5, 127.0, 126.7, 122.9, 121.4, 118.9, 115.5, 55.8, 48.2, 43.2, 31.4, 12.2 ppm.

## Results and discussion

Based on our experience in the construction of diverse heterocycles containing nitrogen via multi-component Povarov reaction (14-16), the preparation of the selected tetrahydroquinoline compounds **5,6** was achieved using the BiCl<sub>3</sub>-catalyzed three-component imino Diels-Alder cycloaddition between toluidine **1**, nitrobenzaldehydes **3,4** and *N*-vinylpyrrolidin-2-one **2** (Figure 1).




Figure 1. Synthesis of nitrophenyl-tetrahydroquinolines using the multi-component imino Diels-Alder reaction.

### 1. Physical description, IR data and yields of the 2-nitrophenyl tetrahydroquinolines (5,6).

Comp.	Molecular Formula	Molecular Weight	IR (KBr), ν, cm <sup>-1</sup>	mp, °C	Yield (%)
5	C <sub>20</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub>	351.40	3394, 2947, 2916, 1666, 1620	222-223	95
6	C <sub>20</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub>	351.40	3271, 2972, 2916, 2854, 1666	242-243	70

These reactions proceeded smoothly in MeCN at room temperature, giving final products that were substances easy to purify and to manipulate. The N-(2-nitrophenyl-1,2,3,4-tetrahydroquinolin-4-yl) pyrrolidin-2-ones **5** and **6** were obtained with good yields: 95% and 70%, respectively (Table 1).

The structures of the C-2 substituted tetrahydroquinolines **5** and **6** were confirmed on the basis of analytical and spectral data and were supported by inverse-detected 2D NMR experiments. IR spectrum characteristic absorption bands of the compound **5** were observed at 3394 and 1666  $\text{cm}^{-1}$ , assignable to the amine and amide groups, respectively, and the nitro group signals at 1512 and 1342  $\text{cm}^{-1}$ . Their mass spectrum showed a molecular ion  $m/z$ : 351 that coincided with the molecular weight (351 g/mol). The  $^1\text{H}$  NMR spectrum of this compound presented the 4-H proton signal at 5.69 ppm, observed as a double doublet with the coupling constants 6.4 Hz and 11.1 Hz. This fact suggested axial-axial and axial-equatorial interactions between 4-H and 3-H protons. On the other hand, the 2-H proton signal was observed at 4.65 ppm with the coupling constants 3.1 Hz and 10.7 Hz that indicated at vicinal axial-axial and axial-equatorial interactions (Figure 2).

The high value of the coupling constant (10.7-11.1 Hz) of the 4-H and 2-H protons confirmed the axial proton

configurations; therefore, substitutes of the C-2 and C-4 positions of the tetrahydroquinoline ring have the equatorial disposition, respectively. On the other hand, it was found by the COSY experiment that the signal at 2.13-1.99 ppm belongs to the 3-H proton, observing the 3-H (4''-H) (2.13–1.99 ppm) and 4.65 ppm (2-H) and 5.69 ppm (4-H) cross peaks interactions (Figure 3).

The nitro-isomer **6** has similar chemical behavior in the spectra data. The chemical structures of the obtained N-(tetrahydroquinolinyl) pyrrolidin-2-one molecules were strongly confirmed through IR,  $^1\text{H}$  and  $^{13}\text{C}$  NMR analyses; however, having a possible mechanism of achieved multi-component condensation, we could anticipate the various diastereomers, *cis* or *trans* configuration. For these reasons, further structural studies were carried out.

### X-Ray Diffraction Single Crystal Study

Samples of both compounds **5** and **6** were grown by slow evaporation in ethanol; however, we could obtain suitable crystals only for the compound **5**. The diffraction data of the compound **5** were collected at 273K using a CCD area detector with graphite-monochromatic Mo  $\text{K}\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ). The data were computed using Bruker-AXS software. For the solution and refinement of the structure, Shelxs-97 (17) and Shelxl-97 (18) were used respectively.






Figure 2.  $^1\text{H}$  NMR spectra of N-[6-methyl-2-(4'-nitrophenyl)-1,2,3,4-tetrahydroquinolin-4-yl] pyrrolidin-2-one (**5**).



**Figure 3.** COSY spectrum of *N*-[6-methyl-2-(4'-nitrophenyl)-1,2,3,4-tetrahydroquinolin-4-yl] pyrrolidin-2-one (**5**).




**Figure 4.** Representation of the unit cell of *N*-[6-methyl-2-(4'-nitrophenyl)-1,2,3,4-tetrahydroquinolin-4-yl] pyrrolidin-2-one (**5**).

Molecular and crystal structures were obtained using Mercury software (19). The molecular structure for the compound is presented in the figure 4. A *cis* conformation of the C-2 and C-4 substitutes is evident, as well as a chair configuration adopted by the tetrahydroquinoline system.


Details of cell data and refinement for the compound **5** are summarized in table 2.

**Table 2. Crystallographic data obtained by four-circle diffractometry.**

Unit cell parameters	$a = 9.109 (2) \text{ \AA}$
	$b = 9.2812 (5) \text{ \AA}$
	$c = 11.011 (3) \text{ \AA}$
	$\alpha = 90.939^\circ (6)$
	$\beta = 100.023^\circ (6)$
	$\gamma = 93.309^\circ (6)$
Volume	913.998 $\text{\AA}^3$
System	Triclinic
Space Group	P-1 (No. 2)
Z	2

**Figure 5.** Molecular packing of the unit cell of N-[6-methyl-2-(4'-nitrophenyl)-1,2,3,4-tetrahydroquinolin-4-yl] pyrrolidin-2-one (5).**Table 3. Atomic positions in the unit cell of N-[6-methyl-2-(4'-nitrophenyl)-1,2,3,4-tetrahydroquinolin-4-yl] pyrrolidin-2-one (5).**

Number	Label	Xfrac	Yfrac	Zfrac
1	O1	0.3653	0.1204	0.8134
2	O2	-0.2751	0.1442	-0.069
3	O3	-0.2013	0.3615	-0.0633
4	N1	0.4385	0.1546	0.3218
5	N2	0.3999	0.2993	0.6826
6	N3	-0.1902	0.242	-0.0246
7	C1	0.2924	0.1252	0.3587
8	C2	0.281	0.2232	0.4685
9	C3	0.4102	0.2053	0.576
10	C4	0.5578	0.2277	0.5329
11	C5	0.5642	0.1906	0.4089
12	C6	0.7043	0.1999	0.3716
13	C7	0.8288	0.2495	0.4491
14	C8	0.8236	0.2933	0.5723
15	C9	0.6885	0.2777	0.6119
16	C10	0.3777	0.2485	0.7914
17	C11	0.3736	0.3751	0.8765
18	C12	0.3898	0.5036	0.8068
19	C13	0.4078	0.4576	0.6777
20	C14	0.1688	0.1496	0.2534
21	C15	0.1696	0.2758	0.1861
22	C16	0.0548	0.3036	0.0958
23	C17	-0.0664	0.2057	0.0701
24	C18	-0.0721	0.0818	0.135
25	C19	0.0456	0.0548	0.2264
26	C20	0.9644	0.3544	0.6587



**Figure 6.** Diffraction profile of *N*-[6-methyl-2-(4'-nitrophenyl)-1,2,3,4-tetrahydroquinolin-4-yl] pyrrolidin-2-one (**5**) simulated in Mercury software.

The packing structure is shown in figure 5 and the powder profile simulated by the single crystal data is shown in figure 6.

Based on the single crystal study of the compound *N*-[6-methyl-2-(4-nitrophenyl)-1,2,3,4-tetrahydroquinolin-4-yl] pyrrolidin-2-one **5** we determined that crystals obtained from ethanol crystallize in the triclinic system with space group P-1 (No 2).

Table 3 shows the atomic positions. Carbon-bound H atom positions were idealized (C-H=0.93 Å), with H atoms riding on the atoms to which they were attached.

## Conclusions

We synthesized two nitro-isomers of *N*-(tetrahydroquinolinyl) pyrrolidin-2-ones using a versatile and simple methodology called the three component imino Diels-Alder cycloaddition. The spectral analysis showed the 2- $H_{axial}$ , 4- $H_{axial}$  configuration; therefore, the di-equatorial disposition of the C-2 and C-4 substitutes confirmed the formation of the endo-adduct during a Diels-Alder cycloaddition process. The

full characterization of the *N*-[6-methyl-2-(4'-nitrophenyl)-1,2,3,4-tetrahydroquinoline-4-yl] pyrrolidin-2-one **5** was possible due to the single crystal X-ray diffraction technique, which provided the following data: compound **5** crystallizes in the triclinic system with  $a = 9.109(2)$  Å,  $b = 9.281(5)$  Å,  $c = 11.011(3)$  Å,  $\alpha = 90.939(6)^\circ$ ,  $\beta = 100.023(6)^\circ$ ,  $\gamma = 93.309(6)^\circ$ ,  $Z = 2$ , space group P-1 [No. 2], and  $V = 1054.0$  Å<sup>3</sup>.

## Acknowledgments

Authors wish to thank *Instituto Colombiano para el Desarrollo de la Ciencia y la Tecnología 'Francisco José de Caldas'* (COLCIENCIAS-CENIVAM), and *Universidad Industrial de Santander*. C.M. Meléndez thanks COLCIENCIAS for the fellowship.

## Financial support

This research was financed by *Instituto Colombiano para el Desarrollo de la Ciencia y la Tecnología 'Francisco José de Caldas'* (COLCIENCIAS-CENIVAM, contract no. 432-2004) and by *Universidad Industrial de Santander* (VIE-UIS, project 5176).

## Conflicts of interest

The authors declare that no conflicts of interest exist in relation to this work.

## References

1. Jones G. Pyridines and their Benzoderivatives: (v). Synthesis, in *Comprehensive Heterocyclic Chemistry*, Katritzky AR, editor. Oxford: Pergamon Press; 1984; p. 3, 395.
2. Katritzky AR, Rachwal S, Rachwal B. Recent Progress in the Synthesis of 1,2,3,4,-Tetrahydroquinolines. *Tetrahedron*. 1996; 52: 15031-70.
3. Kouznetsov V, Palma A, Ewert C, Varlamov A. Some Aspects of reduced Quinoline Chemistry. *Journal of Heterocyclic Chemistry*. 1998; 35: 761-85.
4. Kouznetsov VV, Vargas LY, Meléndez CM. Recent Progress in the Synthesis of Quinolines. *Current Organic Chemistry*. 2005; 9: 141-61.
5. Glushenko TP, Goncharov VI, Aksenov AV. Investigations in 2,3'-Biquinoline Series 24. Synthesis of 3-Hetarylquinolines and their 1,4-Dihydro Derivatives under Conditions of the Vilsmeier Reaction. *Chemistry of Heterocyclic Compounds*. 2008; 44: 973-78.
6. Broch S, Anizon F, Moreau P. First Synthesis of 3,6'- and 3,7'-Biquinoline Derivatives. *Synthesis*. 2008; 2039-44.
7. Chen YX, Yang LW, Li YM, Zhou ZY, Lam KH, Chan AS, Kwong HL. Synthesis of a New Chiral Ligand, 6,6'-Dihydroxy-5,5'-Biquinoline (BIQOL) and its Applications in the Asymmetric Addition of Diethylzinc to Aldehydes. *Chirality*. 2000; 12: 510-13.
8. Ichikawa J, Mori T, Miyazaki H, Wada Y. C-C Bond Formation between Isocyanide and  $\beta,\beta$ -Difluoroalkene Moieties Via Electron Transfer: Fluorinated Quinoline and Biquinoline Syntheses. *Synlett*. 2004; 45, 1219-24.
9. Morsali A, Mahjoub AR, Ramazani A. Zn(II), Cd(II) and Hg(II) Complexes with 2,2'-Biquinoline, Syntheses and X-ray Crystal Structures of  $[\text{Hg}(\text{bq})(\text{SCN})_2]$ . *Journal of Coordination Chemistry*. 2004; 57: 347-52.
10. Trpkovska M, Šoptrajanov B, Pejov L. FTIR Study of the 2,2'-Biquinoline Complex of Dichlorodioxochromium(VI): The  $\text{CrO}_2$  Vibrations. *Journal of Molecular Structure*. 2003; 654: 21-26.
11. Youssef AO, Khalil MH, Ramadan RM, Soliman AA. Molybdenum and Tungsten Complexes of Biquinoline. Crystal Structure of  $\text{W}(\text{CO})_4(2,2'\text{-Biquinoline})$ . *Transition Metal Chemistry*. 2003; 28: 331-35.
12. Pazderski L, Tousek J, Sitkowski J, Kozerski L, Szłachetka E. Experimental and Quantum-Chemical Studies of  $^1\text{H}$ ,  $^{13}\text{C}$  and  $^{15}\text{N}$  NMR Coordination Shifts in Pd(II) and Pt(II) Chloride Complexes with Quinoline, Isoquinoline, and 2,2'-Biquinoline. *Magnetic Resonance in Chemistry*. 2007; 45: 1059-71.
13. Kouznetsov VV. Recent Synthetic Developments in a Powerful Imino Diels-Alder Reaction (Povarov Reaction): Application to the Synthesis of N-polyheterocycles and Related Alkaloids. *Tetrahedron*. 2009; 65: 2721-50.
14. Kouznetsov VV, Mora Cruz U. Transformations of 2-( $\alpha$ -furyl)-4-(2-oxopyrrolinidyl-1)-1,2,3,4-Tetrahydroquinolines, Cycloadducts of the Imino Diels-Alder Reaction: A Simple Synthesis of New Quinoline Derivatives. *Letters in Organic Chemistry*. 2006; 3: 699-702.
15. Kouznetsov VV, Mora Cruz U, Zubkov FI, Nikitina EV. An Efficient Synthesis of isoindolo[2,1-a]Quinoline Derivatives Via Imino Diels-Alder and Intramolecular Diels-Alder with Furan Methodologies. *Synthesis*. 2007; 375-80.
16. Kouznetsov VV, Meléndez CM, Bermúdez JH. Transformations of 2-aryl-4-(2-oxopyrrolinidyl-1)-1,2,3,4-Tetrahydroquinolines, Cycloadducts of the  $\text{BiCl}_3$ -Catalyzed Three Component Povarov Reaction: Oxidation and Reduction Processes towards New Potentially Bioactive 2-arylquinoline Derivatives. *Journal of Heterocyclic Chemistry*. 2010; 47, 1148-52.
17. Sheldrick GM. SHELXS97. University of Göttingen, Germany; 1997.
18. Sheldrick GM. SHELXL97. University of Göttingen; Germany, 1997.
19. Allen FH. Cambridge Structural Database: A Quarter of a Million Crystal Structures and Rising. *Acta Crystallographica Section B*. 2002; B58: 380-88.